

populated States of Massachusetts (0.3 per mille), Pennsylvania (0.4 per mille), Ohio (0.4 per mille), Connecticut (0.5 per mille), New Jersey (0.6 per mille). It is possible that the generally higher level of sanitation in the North, and especially in the more densely populated States, has had much to do in keeping down trachoma, despite what is being brought into the country through recent immigrants, who have settled largely in the northern tier of States.

TABLE 42.—Grand total for trachoma with ratio per 1,000 men.

State.	Number of cases.	Ratio per 1,000.	State.	Number of cases.	Ratio per 1,000.
Oklahoma.....	499	8.21	Montana.....	18	0.66
Kentucky.....	511	7.99	Minnesota.....	43	.60
North Dakota.....	85	4.76	New Jersey.....	44	.58
Missouri.....	270	3.15	South Carolina.....	23	.58
Arkansas.....	130	3.05	Virginia.....	31	.56
Texas.....	290	2.45	Florida.....	13	.54
South Dakota.....	50	2.41	North Carolina.....	28	.51
New Mexico.....	22	2.15	Utah.....	6	.51
Nevada.....	7	2.13	Mississippi.....	19	.50
West Virginia.....	78	2.00	Michigan.....	47	.49
Arizona.....	16	1.93	Connecticut.....	13	.47
Indiana.....	130	1.82	Wisconsin.....	30	.47
Illinois.....	312	1.74	Ohio.....	62	.43
Tennessee.....	104	1.70	Pennsylvania.....	78	.39
Rhode Island.....	20	1.38	Vermont.....	3	.36
Nebraska.....	40	1.36	Massachusetts.....	27	.30
New York.....	314	1.22	Wyoming.....	2	.30
Idaho.....	15	1.20	District of Columbia.....	3	.26
Kansas.....	42	1.09	Delaware.....	1	.20
Colorado.....	20	.89	Maryland.....	6	.16
Alaska.....	1	.82	Georgia.....	9	.14
Alabama.....	42	.81	New Hampshire.....	1	.12
California.....	64	.79	Maine.....	2	.10
Iowa.....	54	.79	State not specified.....	71	.75
Louisiana.....	39	.75			
Washington.....	25	.70			
Oregon.....	11	.69			
			Total.....	3,776	1.37

43. *Amblyopia*.—This condition of “functional” blindness was not detected in a great number of registrants. Its distribution is irregular and probably depended as much upon the training of the medical examiners as upon any biological factors present.

TABLE 43.—Grand total for amblyopia, with ratio per 1,000 men.

State.	Number of cases.	Ratio per 1,000.	State.	Number of cases.	Ratio per 1,000.
Texas.....	248	2.09	Vermont.....	7	0.83
District of Columbia.....	22	1.91	Alaska.....	1	.82
West Virginia.....	71	1.82	California.....	61	.76
Rhode Island.....	24	1.66	Connecticut.....	27	.75
Missouri.....	138	1.61	Florida.....	18	.75
New York.....	397	1.54	Indiana.....	54	.75
Pennsylvania.....	308	1.52	New Jersey.....	56	.74
Maryland.....	54	1.45	Arkansas.....	29	.68
Ohio.....	196	1.37	South Carolina.....	26	.66
Virginia.....	68	1.22	Wyoming.....	4	.60
Oregon.....	19	1.19	Delaware.....	3	.59
North Dakota.....	21	1.18	Montana.....	16	.58
Kansas.....	45	1.17	Tennessee.....	34	.56
Michigan.....	114	1.15	Georgia.....	33	.52
Wisconsin.....	71	1.12	Utah.....	6	.51
Oklahoma.....	62	1.02	Maine.....	10	.50
Nebraska.....	29	.98	Arizona.....	4	.48
Alabama.....	50	.96	Idaho.....	6	.48
South Dakota.....	20	.96	Louisiana.....	25	.48
New Hampshire.....	8	.94	Mississippi.....	16	.42
Massachusetts.....	85	.93	Washington.....	15	.42
Illinois.....	163	.91	Minnesota.....	29	.40
Iowa.....	62	.90	Nevada.....	1	.30
Colorado.....	20	.89	State not specified.....	58	.61
Kentucky.....	57	.89			
New Mexico.....	9	.88			
North Carolina.....	47	.85			
			Total.....	2,947	1.07