2/

1/

Molecular Genetics — the Ideas

Nature of the ideas : easy (means) , not deep : easily related to everyday life ∴ can make models.

school-children can understand.

reason for this : basic theory of chemistry and physics.

(however Helen's ideas (& others today) important in development.)

physics not known is (mainly) irrelevant.

chemistry calculations difficult in practice

Schrodinger book "What is life?" Influence

Its point : Stability, chemical bonds, forces.

Aspects of chemistry & physics used :

Obviously in techniques eg. X-ray diffraction

radioactivity

(ultracentrifuge, electrophoresis, etc)

how especially : bond-distances and angles.

Hydrogen bonds, van der Waal interactions.

Stereochemistry : the one handedness — nucleic acid

— protein,

— sugars etc.

shows shape is important. eg. many enzyme reactions

this why stereochemistry is important.

few basic principles : the helix : equivalence (same point groups)

almost equivalence = coiled coils (viruses)

the same ideas DNA ⇝ RNA ⇝ protein.
their success; except t-RNA.

Watson-Crick idea: replication due to base-pairing

also probably involved in protein synthesis
(though less necessarily more complicated)

the two linear languages: the 4 letter one
the 20 letter one.

related by the code

i.e. only sequence matters; units held together stably by chemical (i.e. strong) bonds.

① how does this give the great variety of structures required?

proteins: the one-dimensional nature of the chemical structure
(remember cross-links)
the three-D nature of the physical structure.

analogy: the structure of a paragraph of English.
(contrast Chinese)

② why two languages:

the importance of neutral selection — but basic idea in all biology.
(Haldane: random mutations)

3

the requirements for natural selection

① geometrical replication — fairly exact

② meaningful random mutations necessary
i.e. "mistakes" — "rare accidents"
Natural selection. | gives opportunity to improve.

③ "versatility"

[④ requirement for an "association" i.e. a cell]
Gene + products must stay associated.

Thus nucleic acid: from 2 (+3) stranded structure.
idea for replication, but very
inversatile. (probably no Watson-Crick pair
possible)

protein

protein: main structure is α-helix
single-stranded. marginally stable in
water (hydrophobic group rich inside)
20 or so diff residues, with a variety of
possible chemical function.
But probably bad for replication.

Hard not clear that an electronic system
could not do it. May be due to "origin" i.e. historical accident.
rather than coupling system is better

Are two distinct languages necessary a theoretical
grounds i.e. to avoid confusion between the master copy (the gene)
and the working tools (many copies required) ~~etc~~ ?

novice process is unidirectional explain

the central dogma (explain)
irreversibility due to energy put into it.
(also helps to increase accuracy !)

partly explains why ~~inherited~~ acquired characteristics
not inherited.

---

maths in biology ~~&~~ for special purposes

e.g. x-ray diffraction

is any theory too difficult for experimentalists? } theory of unwinding helices (the pure theoretician)
really, the economics of the cell.

lack of application of information theory in
molecular genetics yet.

probably due to very low noise of the channel.
in fact mistakes are too low
(natural selection not operated to reduce them)
repair mechanisms!

more important overall conclusion.

in ~~Biology~~ Genetics individual atoms matter, even only in

Special places.
for many years
obvious that genetic interaction at or near atomic level.

example of Sperm : size. half interaction :

human complexity : there ---
how does atom produce its effect?
the Sickle cell case

explain diseases (two genes) 1 aa. change.

then 1 (x2) "atoms" wrong. how does this kill.

fertilized egg. cell division : many 10^{12-13} cells.

cells in same-person : each many copies of

(stable) m-RNA : each molec many, many copies of protein.

10^6 cells / second
(life 3 months?)

~~x 6~~ 10^7 molecs life?

Molecular Genetics

Private { Stress the genetic message is
a linear one, its units a c polymer
then i a lot of it; and its units at
the atomic level.

Problem in the future : control : embryology (shape + form)
origins. brain.

Remarks on methods

Some experiments done on intact cells; some on purified parts of broken cells; some are both techniques.

methods must (So can be repeated many times) be rapid, and sensitive i.e. work on small quantities

techniques for Separation: ultra centrifuge
electrophoresis
Chromatography + columns
— paper.
(counter current)

Nuclear physical-chemistry.

techniques for detection and recognition
— U.V. absorption
— radioactivity.

Subject has advanced much because methods so powerful (in addition to a good active theory)

1/

The Structure of DNA by X-rays

work of Wilkins + Franklin

Nature of Structure — S

(also calf-thymus)

preparation of DNA : break open cells / : selective precipitation.

can be got fairly pure, but "damaged" fibrous mass.

technique of pulling fibres — to get alignment. an art

control of humidity (also the salt, and pH)

taking x-ray photograph. Longish exposures. need fair amount of material (contrast e/m)

what do x-rays see? electron density : not base sequence.

repeating structure gives discrete "spots". Examples : S

note a fibre diagram; crystallites, etc. S

To solve structure: Fourier transform : amplitudes but not phase

(pseudo-echo symmetric). Transform of a helix. ~~bases~~ stacks of bases

backbone S

Model building : distances and angles known.

"rotating" and "physical contact" are not known, but obey rules.

Good for for all models

S

base pair in a crystal. ethyl guanine | other crystals.

methyl cytosine.

[AT pair different]

Meselson - Stahl expt

ultra centrifuge (commercially available) - two types) swinging bucket

various ways to use : boundary sedimentation.

Sucrose gradient "

density gradient centrifugation (Vinograd)

explain : method.

use of heavy isotopes. ie pair N_{14} + N_{15}

describe experiment : grow many generations on N_{15}

time zero : change to N_{14}.

later time, break open + put into cell.

DNA "chromosome" broken into pieces, but still high MW.

later do not expect S

Results S

Show a duplex.

Subsidiary experiments show not end to end

and almost certainly a two chain duplex, not a four.

3

KORNBERG — JOSSE

~~Aim~~ · Nearest neighbour experiment. use of radioactive isotopes

diff. DNAs have diff average composition.

Sequence not random.

design of experiment. ——— S

result. ——— S
(control — can be done twice)

shows chains run in opposite directions..

4

Fine Genetic Mapping

Results by genetic methods show

ⓐ genes arranged in a linear (circular) array i.e. not 2D

ⓑ units of a gene in linear

ⓒ many distinct sites inside a gene.

BENZER

my example: the r_{II} locus of phage T4. explain growth cycle

r_{II} mutants.

"Spontaneous"

chemical – base analogues

[acridines]

	B	K
mutant	grow but look different	don't grow
wild	grow	grow.

How ⓐ can we pick up mutants

ⓑ selection technique

method of genetic cross: using selection technique.

map distance.

deletion mapping.

result map unit many sites. ——·—— S

If lots of time do cis–trans test.

to show two genes, or cistrons.

Colinearity of gene and protein — State result

Yanofsky's results : A protein of tryptophane synthetase of E. Coli.

Work of Sarabhai, Stretton & Brenner, and Bolle.

Head protein of phage T4.

major protein of ~~head~~ phage. major late protein.

∴ most radioactive label goes into this protein

∴ no need to purify.

Special class of mutants "amber"

	Host A	Host B
Amber mutant.	mutant chain terminated	"wild" chain finished.

Special gene & which does this

Then can be grown in Host B
but tested in host A.

Series of amber mutant, picked up by hand work. map — S

~~Host~~

Chemical test in host A.

Label in vivo all with one radioactive amino acid

Break open. (remove DNA) digested with trypsin

do one-D paper electrophoresis to see which peptides there. Repeat with ① diff. amino acids for ② chymotrypsin ③ diff. amber. Result — S