

Randall (B. A.) Dr. J. D. Billings, M.D.
with the compliments
of the writer.

The Refraction of the Human Eye.

A Critical Study of the Statistics obtained by
Examinations of the Refraction, especially among
School Children.

BY


B. ALEX. RANDALL, A. M., M. D.,

Ophthalmic and Aural Surgeon to the Episcopal and the Children's Hospitals, Philadelphia.

FROM

THE AMERICAN JOURNAL OF THE MEDICAL SCIENCES.

JULY, 1885.




THE REFRACTION OF THE HUMAN EYE.

A CRITICAL STUDY OF THE STATISTICS OBTAINED BY EXAMINATIONS OF THE REFRACTION, ESPECIALLY AMONG SCHOOL CHILDREN.

BY B. ALEX. RANDALL, A.M., M.D.,

Ophthalmic and Aural Surgeon to the Episcopal and the Children's Hospitals,
Philadelphia.



THE great importance of full sharpness of vision is a matter which is steadily obtaining wider-spread recognition, and is demanding still more earnest consideration of all thinking persons. Defective sight is growing more notable for a variety of reasons, and the means of amending it are progressively improving and are more widely employed, thus furnishing constant reminders of the need of such help; yet they are making only the more distinct those classes of cases which are but partially, if at all, remediable, and driving us to the more thorough study of the causes of these defects and the means for their prevention. The progress which has been made in all branches of knowledge, especially the natural sciences, is constantly increasing the field to be covered in the usual educational process. The demands made upon children at their lessons have steadily increased; while the assumption of the matter of education by the state, as a duty owing to itself and its citizens, has carried the influence of the schools, sometimes by compulsion, into every rank of life. Conditions, therefore, which not long since invested only the few, are now the environment of the community. Census reports point to a steady and rapid decrease, in every country making claim to civilization, of that portion of its inhabitants unable to read and write; while means and attractions to the exercise of these powers are offered by veritable floods of literary productions and reproductions, by postal facilities which would have been deemed a few decades since utterly beyond belief, and by an immense broadening, for almost every one, of the field of every-day interest. Some information on almost all points is expected of every one; and the most abstruse dis-

coveries of modern science in revealing new worlds, measuring the globe, or investigating its components and the creatures which live or have lived upon it, are quickly absorbed into the fund of general information, and go to widen or to modify that stock of knowledge of which all are expected to be cognizant.

And the means to the attainment of the major part of all this is the eye. Most children begin to study the alphabet and its combinations before the fifth year, and a few years more finds them all readers. School tasks surround and occupy them much of the time, and of what remains not a little is given to amusements hardly less exacting in their demands for the use of the eyes in accurate vision. The general spread of the employment of artificial illumination has lengthened greatly the hours during which the eyes can be rigorously exercised, and the child whose school work is finished turns to other reading for recreation. In the general walks of life vision, and good vision at that, is being more and more strictly required—how much we are hardly likely to appreciate at first. Yet if we think of any one, of whatever age or rank, suddenly deprived of vision, we will, perhaps, in some measure realize how essential to almost every employment is the lacking sense. How few are the paths now open to the blind, and in them, how difficult it is that they should at all cope with seeing competitors! The disadvantages of those with defective vision differ in degree rather than in kind. In every one of our eye clinics the cases are numerous where persons reject with alarm the advice that they should wear glasses, knowing, as they do, that such an assumed confession of visual defect may cost them a present situation, and will certainly count to their prejudice in seeking any new place. All this is but a fraction of the evidence showing how much is required, even outside of the schools, of the visual powers of those who have them in the usual measure.

The present century has seen the introduction of methods and instruments of precision into almost every branch of study, most notably in all which would lay any claim to scientific value. The inductive method is almost universal, and each advance is a generalization from extended observations. The observer works for the instruction of the community as well as himself; his results must receive confirmation in order to meet general acceptance; and to this end standards of comparison more or less constant and methods free as far as possible from the personal equation have become essentials. Advancing with the general realization of the importance of good vision, ophthalmology has furnished means more and more exact for the determination of the visual acuity and for the measurement and correction of its defects; until we are now in position to obtain data in these matters as strictly accurate as any in the whole range of natural science. In the mean time, investigation after investigation has been made in the

matters of vision and refraction, embracing enormous numbers of individuals, in the endeavor to answer the numerous hygienic and social questions involved in the subject of defective vision. Early realizing that, to obtain results of general applicability, eyes presumably normal must be the objects of study, and that in youth many of the causes of defect have not yet come into play, the eyes of school children have been most frequently examined, both to obtain information as to the important subject of the special hygiene of schools, and also to gain data as to what is to be regarded as the standard eye, the typical form, the normal, which the various conditions of imperfection have failed to attain. This standard determined, and the statistics of the primary variations from it once settled, we will be in position to study the questions of the general prevalence of defective vision, the causes which are effective in evoking it, and the measures to be employed in combating the evil.

The foundations of such a wide-reaching study must, of course, be broad. Furthermore, they must be sure; lest the crumbling of some of the constituents should imperil the stability of all that is built upon them. The first condition would seem well answered, since no student of the subject, not even Prof. Cohn, has up to the present time had the patience to collect all the statistics of the matter already in print; yet arrays of scores of investigations, based upon thousands of individuals, are to be found in the summaries which have been given by various authorities. In the other direction much remains to be desired in not a few of these studies, so much, indeed, that some of them can be employed in any summarizing of results only within the narrowest limits. Setting out in many instances with some single object in view, such as the question of the prevalence of myopia, attention has been given to it alone, and their negative showings in other directions are often valueless, even where the positive results are reliable. Yet they are often grouped as all equal and competent portions of evidence, and advantage is taken of their wide divergence on many important points, either to employ the convenient ones in the support of any fallacy, or else, by holding them up into view in their apparent contradiction, to throw discredit upon all such investigations and the so-called science which they in some measure represent. There has been almost no attempt to combine a critical examination of the methods, details, and results of this extensive series of investigations, with a summary embracing all that are accessible. No light work nor a pleasant one, dealing as it must with comparisons more or less odious, yet a task which the writer here undertakes in the belief that valuable results are thus to be attained. That these results will display any striking novelty is not to be hoped; but that they may give added weight to concurrent testimony derived from other sources, and perhaps widen the acceptance of views not yet receiving their due appreciation, seems abundant

reason for the labor necessary to educe them. Where a hundred investigators have expended years of study in gathering statistics on so weighty a subject, surely there is place for one to combine the data thus gathered. If only to show that much of the labor has been wasted in the past, and to give indication of the points to-day demanding elucidation and the methods by which they are to be studied out, this study would appear to be called for. That it can give or disseminate much other valuable information will, it is hoped, be made to appear.

A comparison of the numerous investigations which have been made in the field before us shows the utmost variation in their aims, methods, and results. Great as are the differences, however, the studies will repay careful scrutiny, and in few instances will we fail to find reason for discrepancies. In most of the examinations the question of the occurrence of myopia and its increase was paramount; in many, it alone was studied. Indeed, the earliest investigators knew of nothing else in our field to seek, and the methods at their command were hardly sufficient for its discovery. Even after hypermetropia and astigmatism became well known they were rarely sought with any care, and vision equal to normal was considered, sometimes even defined, as synonymous with emmetropia. In many of the investigations each eye was not separately tested, so that practically only the better eye was studied, and any difference between the two overlooked. Investigating the subject with rather vague hygienic views, the individual was quite naturally the centre of interest. If he could see with normal sharpness, whether with one eye or both, he was not a myope, and the requirements of the matter were considered to be satisfied. Among the later examinations such oversights are, naturally, rarer; yet often the meagre reports available either give no information on such points, or afford it in such indefinite terms as to leave them still in doubt.

In many of the studies the nature of the refraction has been carefully determined; but the degrees of the anomalies found have either not been determined or not reported. Others state these important details for the class, the age or the year of school life; yet, omitting the mention of the actual numbers of individuals or eyes embraced under each heading, the results, valuable as they may be in themselves, cannot be combined with those of others. In some of the cases where the ophthalmoscope was employed in the examination of every eye (and in the many where it was not used, a most important aid to accuracy was omitted), it was used to seek for lesions and not to measure the refraction. It is needless here to repeat that in the aims held by some of the investigators such details had no place; and that in studies so comprehensive in the matter of numbers examined, the introduction of the methods of extreme refinement would have been impracticable. What most concerns us is the stability, the correctness, of our foundations. The subject here taken up is the information which these examinations *can* afford as to the usual refraction of the

human eye; in health, as far as we may exclude pathological conditions in youth and in age, if the data are to be found: to extract all that is germane to our purpose. And when grave omissions or vexatious errors, generally typographical, have to be pointed out, it will be with full realization from the writer's own experience, of how hard it is to avoid error in statistical reports, and how impossible for one man to comprehend in his investigation all those points that to another may seem most interesting or essential. And if, in passing, we cannot conceal a regret that some study, most valuable in some respects, gives us no information in others, it will be with no thought of adverse criticism—rather in testimony to the value of that which is given.

To begin this study with a series of definitions, where the subject has been so much written about, may seem unnecessary, if not presumptuous. Yet, since loose employment of technical terms constitutes the principal ground of fallacy in statistics, the cardinal vice, which can make figures the most lying of witnesses, it is essential that some points be absolutely defined, and then the data of various observers be marshalled under these headings with the strictest accuracy which the details at hand will permit.

Waiving technicalities as far as possible: the *refraction* of the eye is the relation of the situation of the percipient retina to the focus of the lens system of the passive eye, for rays of light coming from distant objects. When, without any effort, the focus for such rays falls on the perceiving layer of the retina, the eye is *emmetropic*. When the eye is relatively too long to meet this condition, it is *myopic*; its retina is in position to receive the focus only of rays coming from some point not remote; it is *nearsighted*. When relatively too short, the eye is *hypermetropic*; without accommodative effort, or its equivalent, the eye can focus no rays of light that occur in nature; distant objects form undefined images upon the retina; near objects form images still less clear.

Vision is the perception of external objects through the agency of the rays of light coming from them and impressing the retina. Its acuity depends upon the formation of a sharp image upon this membrane, and upon the capacity of it, and the visual centres behind it, to take cognizance of the image so found. Granted the latter conditions in full measure, the acuity of vision depends upon the transparency of the ocular media and the refraction, primary or fictitious, of the eye. By fictitious refraction is meant every deviation from its primary condition above defined, whether produced by accommodative effort or by optical appliances, such as lenses. Conditions of imperfect retinal perception and of non-transparency of the media before it, may for our purposes be grouped under the heading *amblyopia*, and practically laid aside, and we have before us the subject of the visual acuity of the human eye as dependent upon its refraction, this last being divisible into real or primary, and fictitious or apparent, refraction. The real or primary refraction is the

object of our present search; and we will endeavor to winnow out of the mass of results furnished by a long series of investigations those which have bearing upon it, and try to solve with their aid the question—What is the normal refraction of the human eye?

Since we have at our command easy methods of determining with the utmost accuracy the visual acuity in any case; this, the most salient practical point of our study, can be directly measured; and as by wide observation we have obtained reliable data as to the average sharpness of vision, we can make ready use of it as a means to our end, only noting that the tests must be strictly applied. The farthest point of distinct vision for the myopic eye is not remote, hence normal vision at a distance without lenses at once excludes myopia: but the opposite condition of hypermetropia is far more difficult to exclude. Exercise of the accommodative power, which in the myope only makes distant vision worse, can in the hypermetropic eye cover up the defect of refraction and raise vision to the normal; nay, more, the small pupil, and probably the tension of accommodation and its effects, may raise vision above that which we hold to be the average. Vision above the normal is rather indicative of hypermetropia. The accommodation must be set aside in order to reveal the real refraction, above defined. A convex lens held before the eye will render the rays coming from distant objects convergent, and only a hypermetropic eye can focus such rays, and see normally at a distance through a convex lens. The accommodation used to render the eye apparently emmetropic is now unnecessary, and can only impair vision; it may, therefore, be laid aside, and the eye then *manifests* its hypermetropia. But a lifelong habit will not always yield in an instant. The accommodative tension may be maintained, distant vision is made worse by a convex lens, even the weakest; and we are driven to other methods to attain our aim. Dark surroundings are conducive to relaxation of accommodation, and the glare of light reflected into the eye by the ophthalmoscopic mirror may aid, or at least not counteract, the influence. This method, then, in expert hands, may as it were surprise the eye into a revelation of its defect. Yet all such measures may fail, and nothing short of absolute paralysis of accommodation by a mydriatic will tell the whole story. Irritative causes have perhaps set up a cramp of the ciliary muscle, a spasm of accommodation, which will yield but obstinately to the most powerful mydriatic.

The accommodation, then, is the factor which introduces uncertainty into the investigation; study of it can sometimes clear away doubts. The range of accommodation is at each age approximately a constant: if a part of it be called into play to cover up hypermetropia, by thus much will the range be shortened. Here we have not only a clue to the existence of such a condition, but also some indication of the amount of accommodation thus rendered *latent*; and hence of the amount of hypermetropia

neutralized. Where astigmatism is absent, very accurate results may often be thus obtained. Similar measures may be employed to determine the presence and also the degree of myopia, and to exclude the cases where nearsight is simulated by accommodative spasm; but the same difficulties may be met and nothing less than total paralysis of accommodation will enable us to claim absolute accuracy. Still more is this the case with astigmatism; and although we can in addition to other methods, get absolute results by ophthalmometric measurements as to corneal asymmetry, we have only one—the most important—of the series of factors involved.

To turn, now, to the practical workings of an examination, let us see what information we can obtain, and how we may prove the existence of emmetropia, or other conditions of refraction. Testing of the visual acuity will show it to be normal or subnormal—the former excluding myopia. If subnormal, the use of convex or concave lenses may show the presence and degree of *absolute hypermetropia* or *apparent myopia*, respectively. In all these tests each eye must be separately tried; but sometimes the testing of both together may be also advantageously employed. Convex glasses held before eyes with normal vision may reveal hypermetropia (“H. facultative manifest” of Donders), or in case of their rejection will speak for apparent emmetropia. The various tests for astigmatism may show the presence of this defect, and indicate, perhaps correctly, the meridians of greatest and least refraction in the media of the eye. These data may be confirmed by the ophthalmoscopic examination, most especially in the direct method and in retinoscopy; the cornea may be measured with the ophthalmometer, and the range of the accommodation may be determined and compared with the standard furnished by Donders. Concurrence of these various tests will greatly enhance the confidence which would be due to the findings of any one of them alone; and the presumption will be strong that eyes, which by all tests appear emmetropic, are so in fact. Only by paralysis of the accommodation, however, can this be absolutely settled, and the emmetropic eye be proven to be actually such; then no latency of hypermetropia could enlarge the list of seeming emmetropes, and no spasm of the accommodation could decrease it by causing apparent myopia. Then, too, the *degree* of any error of refraction could be exactly measured—all previous tests claiming to furnish but an approximation.

One principle must rule in all these tests. Accommodation is to be done away if possible. It can increase temporarily the refraction of the eye, lessening or covering hypermetropia, increasing the degree of myopia, or making apparent myopes of those really emmetropic or hypermetropic. There is nothing tending to the contrary—no counter-force to oppose it or to introduce error in the other direction. The lowest grade of myopia, therefore, or the highest grade of hypermetropia, obtainable by any method of competent accuracy, is the nearest approximation to the real refraction. All data in this direction are positive; all to the contrary, merely nega-

tive. Hypermetropia proven by any one test, for example, is to be considered as the true refraction of the eye, although every other test points to the contrary. The one test, to stand thus in conflict with all others, must be well based; it should be repeated to obtain confirmation from itself; but standing still, it is to be accepted as correct—all the others are probably vitiated by the deceiving activity of the accommodation. Of the various methods of examining the eye with unparalyzed accommodation, that by the ophthalmoscope is most frequently able to act such a part. The conditions of accuracy must be fulfilled and the observer must be competent; then its findings can be maintained against any array of others, which point towards myopia, *i. e.*, in the direction in which accommodative activity would tend.

In the abstracts which follow, the original reports of the examinations have been strictly followed in every case where they have been accessible; the source of information actually employed being given in the appended bibliography, to which the numbers refer. Abstracts and reviews have also been studied to obtain all possible collateral information; in this regard, and also as a source of most quotations which have had to be made at second-hand, use having been most frequently made of the "Jahresbericht über die Leistungen und Fortschritte im Gebiete der Ophthalmologie." Our study being of *eyes* and not individuals, this as well as convenience or conciseness has here and there dictated a recasting of the forms in which the results have been given; but effort has been made to do the least violence consistent with our needs to the observer's own arrangement, and none whatever to his actual results. Occasionally, palpable error in the summing up given by the examiner has driven us to work over his detailed results, and figures have been thus obtained differing (rarely in notable degree) from his totals. Information as to the actual *numbers* being more important to our ends than percentages, these numbers have been worked out when not given—always with the proviso that they are merely approximations. All results which are not in perfect accord with those of the investigator cited are indicated by parentheses, or, if derived by the writer from percentages, etc., by brackets. The varying nomenclature and standard of the schools in which many of the investigations have been made, and of their classes, makes uniform arrangement in this regard well-nigh impossible. The designation of the equivalent American school has been sometimes given, but the untranslatable "Realschule" and other similar terms, have been retained. Age, being looked upon as the best basis of comparison for our purposes, has been so employed whenever the data permitted. As to the "Gymnasium," it may be remarked that its curriculum extends generally from the fourth year of school-life about as far as the Sophomore studies of our American colleges; while in the "Realschule" the course is approximately parallel, but with lighter stress upon the classical studies.

No. Date.	Examiner.	Character of examined.	Age.	Persons. Eyes.	Em. per ct.	M. per ct.	H.m. per ct.	H. per ct.	As. and Amb.
1 1813	Ware	Foot guards, England	10,000	0=0?			
" "	" " " " " "	Chelsea Military scholars, England	1,300	3 0 25.7			
" "	" " " " " "	University students, Oxford	127	32 25.2.7			
2 1848	Szokalski	Primary scholars, Paris	6 300	0 0 ?			
3 1866	Schürmayer	College and Lyceum pupils, Paris	877+	114 [11.6]			
4 1861	v. Jaeger	Grammar-school pupils, Baden	930	46 4.9			
5 1866	Kuete	Normal school, etc., pupils, Baden	2172	392 18.4			
6 1867	Roek	Normal children and soldiers, Vienna 9d. 257.	(800) 700	223=31.8	416 59.4		61=8.7?	
7 "	" " " " " "	Free charity school pupils, Leipzig	213 (251.4) 1	[12]	48 [32.5]			
8 "	Alexander	School-children, Brunswick	2250	?			
9 "	Gaertner	Primary scholars, Aix	869	231 83.7			Am. 34
" "	" " " " " "	Theological graduates, Tübingen	+19	138	41=14.9	22 1.4	?		
" "	" " " " " "	Village primary pupils, Breslau, etc.	6-13	1486	1408 94.7	32 1.4	16= 1.1		
" "	" " " " " "	City elementary pupils, Breslau	6-14	4078	4245 85.3	334 6.7	127 2.5		
" "	" " " " " "	Secondary and Grammar pupils, Bres.	7-17	1260	595 78.9	109 8.65	52 4.12		
" "	" " " " " "	Realschule and Gymnasium, Breslau	9-22	2336	1682 72.	539 23.	44 1.9		
10 1868	" " " " " "	University students, Breslau	17-27	410	134 32.7	244 59.5	15 3.7		
11 "	" " " " " "	Type-setters, Breslau	20-71	132	51 38.7	68 51.5	10 7.6		
12 "	Thilenius	University scholars, Rostock	314	175 55.7	96 30.6	9 2.9		
13 1870	Schultz	Gymnasium scholars, Upsala	12-16	431	?	159 36.9	?		
14 1871	Erismann	Pupils of various schools, St. Petersburg	8-24	4338	1132 26.	1317 30.2	1889 43.3		
" "	" " " " " "	Village scholars, Silesia	6-13	240	14 3.	4 0.9	370 77.		24
16 "	Makiakoff	School children, Russia	769	482 63.5	251 33.5	23 3.		
17 1872	Cohn	Gymnasium pupils, Breslau	6-22	361	187 52.	125 35.	25 7.		24
18 "	Krüger	Gymnasium pupils, Frankfurt	9-19	203	159 39.	162 40.	(72 17.7)		
19 1873	v. Hoffmann	Gymnasium pupils, Wiesbaden	9-19	256	140 27.3	157 36.5	[101 19.7]		
" "	" " " " " "	Girls prep. & gram. pupils, Wiesbaden	6-15	403	203 50.4	83 20.6	87 21.5		9
" "	" " " " " "	Boys prep. & gram. pupils, Wiesbaden	6-16	568	321 56.5	67 12.	37 6.5		23
20 "	" " " " " "	Gymnasium pupils (1872), Vienna	409	243 29.7	375 45.8	158 19.3		35
" "	" " " " " "	Gymnasium pupils (1873), Vienna	389	144 37.	108 42.7	69 17.7		4
21 1874	Ott & Ritzmann	Gymnasium pupils, Schaffhausen	122	124 51.	98 39.3	24 10.		
22 "	Ott	Realschule pupils, Schaffhausen	12-14?	164	324 146 45.	37 11.1	122 37.7		
23 "	" " " " " "	Old villagers, Silesia	60-84	100	200	25 12.5	114 57.		As. 19
24 "	Gayat	Pupils of various schools, Lyons	6-14	690	440 73.3	52 8.7	108 18.	114 57.	12?
" "	" " " " " "	Girls Grammar school, Munich	10-17	179	[72] 40.	88 49.2	19 10.7		
25 "	" " " " " "	Realschule pupils, Berne	10-18	430	279 64.9	125 29.	18 4.18		
26 "	" " " " " "	Negro prim. & grammar school, N. York	5-19	457	159 17.7?	12 2.6	239 67.		360 76.5
27 1875	Callan	Gymnasium pupils, Vienna	480	187 33.1	177 34.6	49 10.5		132 28.3
28 "	v. Reuss	Gymnasium pupils, Vienna	480	187 33.1	177 34.6	49 10.5		

1 Only those having eye-trouble were sent for examination.

(Atropine)
438=91.2

?
or > 1. D.

No. Date.	Examiner.	Character of examined.	Age.	Persons. Eyes.	Em. per ct.	M. per ct.	H. m. per ct.	H. per ct.	As. and Amb.
29 1875	Pfugger	Teacher recruits, Switzerland	20-25	529	3036	112=21.2	337=11.76	1441=47.47	
30 1876	Courad	Gymnasium pupils, Konigsberg	6-20	1618	909=29.94	678 22.33	126 63.		
31 "	Adler	Deaf-mute pupils, Vienna	8-19	200	64 32.	10 5.			
32 "	Spalding	Primary and Grammar pupils, Portland	5-15	2372	244 11.7	244 11.7			
33 "	Pfugger	Pupils of various schools, Lucerne	7-23	1846	862 46.	184 10.	800 44.		
34 "	Emmert	Pupils of various schools, Berne, etc.	6-55	2145	422 9.82	542 12.61	3308 77.		
35 "	Loring & Derby, R. H.	Pupils of various schools, New York.	6-21	2365	1531 67.59	448 19.78	286 12.63	?	
36 "	Schelding	Gymnasium pupils, Erlangen	8-	746	187 25.1	371 49.7	188 25.2		
37 "	Koppe	Pupils of various schools, Dorpat	661	[166 12.4]	[224 18.8]	687 61.5	[904 67.8]	As. 92
38 1877	Kotlmann	Normal scholars, Hamburg	8-20	413	279 20.9	270 20.3	8 2.		
39 "	"	Realschule pupils, Hamburg	232	240 58.	157 38.			
40 "	"	Grammar scholars, Hamburg	310	[80] 25.			
41 "	"	Pupils of girls private school, Hamburg	104	[77] 25.			
42 "	"	Pupils of girls private school, Hamburg	218	[18] 17.			
43 "	"	Girls' normal school pupils, Hamburg	45	[48] 22.			
44 "	"	Primary scholars, Hamburg	296	[19] 42.			
45 "	Bacon	Realschule pupils, Hamburg	402	[36] 12.	34 5.5		
46 "	Becker	District scholars, Hartford	7-17	308	464 75.3	102 16.5			
47 "	"	District scholars, Heidelberg	287	[120] 18.			
48 "	"	Gymnasium pupils, Heidelberg	616	154 53.	100 35.	33 12.		
49 "	"	Grammar pupils, Cincinnati	291	[34] 13.			
50 "	"	Primary district pupils, Cincinnati	630	507 80.5	83 13.5	37 6.		
51 "	"	College students, New York	549	651 59.	383 30.3	105 9.6		
52 "	"	Tech. Institute pupils, Brooklyn	30 0	372 62.	79 12.5	126 21.		
53 "	"	College freshmen (Harvard) Boston	122	[36] 29.5			
54 "	"	Gymnasium pupils, Magdeburg	325	[156] 49.			
55 "	"	Grammar school pupils, Magdeburg	388	[171] 44.			
56 "	"	Watchmakers, jewelers, lithogra- phers, Breslau.	19-71	172	28 16.2	25 14.5	As. 3
57 "	"	Lycenum pupils, Lyons	5-23	1016	116 67.4	568 28.	= or > 1 D.		
58 "	"	Gymnasium pupils, 1876, Schaffhausen	120	1300 64.	98 40.8	87 4.2		
59 "	"	Recruits, grad. of Gynn., etc., Munich.	18-24	284	90 37.5	161 58.	62 21.6		
60 "	"	Pupils of various schools, Caucasus	1259	[529] 42.7	[365] 29.7		[428] 34.7	= 5.
61 "	"	Gymnasium pupils, I and II, Berlin	61	122	76 62.3			
62 "	"	Med. Students and graduates, N. York	21-31	14	3 22.	0 0	11 78.	
63 "	"	Gymnasium pupils (II & I), Hesse	177	354	[209] 59.3	202 23.7		
64 "	"	District school, boys, Dresden	6-15	432	470 55.	182 21.3	238 31.2		
65 "	"	District school, girls, Dresden	6-14	382	317 41.5	509 27.3	121 16.4		
66 "	"	City school, boys, Dresden	6-14	368	736	429 58.3	186 25.3		
67 "	"	City school, girls, Dresden	6-14	364	723	409 56.3	169 22.9		
68 "	"	Girls school pupils, Dresden.	7-16	171	337	[187] 55.5	55 16.3		

3 Atropine.

3 The percentages foot up 110 per cent. through a typographical error.

1 Perhaps the same as No. 112.

No.	Date.	Examiner.	Character of examined.	Age.	Persons. Eyes.	Em. per ct.	M. per ct.	H. per ct.	H. per ct.	As. and Amb.
1878		Haeuel	Realschule, Dresden	8-17	209	417	[234=48.5	161=38.5	52=12.4]	
"	"	"	Kreuzschule, Dresden	9-21	554	1100	460 41.5	654 50.	86 7.8]	
"	"	"	Gymnasium, Dresden	9-21	480	953	450 46.9	449 46.8	64 5.62]	
"	"	"	Royal Seminary, Dresden	13-21	218	432	164 37.9	236 54.8	32 7.4]	
"	"	"	Private Seminary, Dresden	13-20	131	291	103 39.5	134 51.3	24 9.2]	
56	1879	Derby, Hasket	College freshmen, Amherst	15-24	321		145 45.2	114 35.9	67 20.9	
57	"	Horstmann	Young children [Berlin?]	8-24m.	57	96	13 13.5	9 9.4		(Atropine) 74=77.
58	"	Netolicka (vide 82)	Pupils of various schools, Graz	1229	2458	(1426 57.7)	735 29.9	234 9.3	[1660 67.7]
59	"	Jest	Pupils of various schools, Zittau	1717	3434	622 15.5		
60	"	Nenti	Primary and lyceum pupils, Marselles	26	51	14 28.	0 0.	85 71.4	(37)
61	"	Kotelmann	Uncivilized, Hamburg	3m-46y	283	566	182 31.62	110 19.43	273 48.83	
62	"	Kotelmann	Gymnasium pupils, Wandsbeck	8-17	1636		86 5.3		
63	1880	Smith, Priestley	Pupils of Board-schools, Birmingham	7-13	357		72 20.		
"	"	"	Pupils of normal schools, Birmingham	18-23					
64	"	Ely	Infants, New York	1-60d.	111	154	21 14.	27 18.		(Atropine) 106 69. (Atropine) 28 70. (133 11.7)
65	"	Horstmann	Infants, [Berlin?]	1-20d.	40		8 20.	4 10.		
66	"	Dennett	Pupils of various schools, Hyde Park	4-20	1133		894 75.3	95 8.4	133 11.7	
67	"	Behelm-Schwarzbach	Pupils of various schools, Australia	8-17	3397		298 7.9		
68	"	"	Maori children, Australia	10-15	104		15 14.4	3 2.9	86 82.6	
69	"	Emmert	Gymnasium pupils, Berne	6-18	203	406	36 8.9	52 12.8	318 78.3	
70	"	Florschütz	Pupils of var. schools (1873), Coburg	6-20	2041		1329 65.1	434 21.3	72 3.5	
"	"	"	Pupils of var. schools (1877), Coburg	6-20	2323		341 14.7		
71	"	Herzenstein	Military Gymnasium pupils, Orel	6-14	472		242 51.2	69 14.6	?	
72	"	Schillbach	Gymnasium pupils, Jena	[206]		21.6-66.6		
73	"	Netolicka (vide 82)	Pupils of primary schools, etc., Graz	<10d.	281	562	10 2.	0 0.	552 98.
74	1881	Königsfeld	Medical students, Vienna	108	216	81 37.5	116 54.4	
75	"	Cohn	Common school pupils, Vienna	6-14	472		242 51.2	69 14.6	161 34.1	
76	"	V. Reuss	Higher schools, Darmstadt	9-21	1389	2778	1767 42.7	801 34.27	252 8.3	824+
77	"	Weber	Pupils of various schools, Philadelphia	6-21	1212	2422	2177 11.2	332 13.7	1792 74.1
78	"	Risley	Dea-mute pupils, Bordeaux	200		15 7.5	24 12(+)	As. (1330)
79	"	Badal	University students, Utrecht	17-38	410	820	389 47.4	286 35.	198 19.2	As. (105)
80	"	Collard	Gymnasium pupils, Schletz	11-21	149	298	133 44.5	149 56.	16 6.3
81	"	Westphal	Keatschane pupils, Graz	254		136 53.5	75 29.5
82	"	Netolicka	Pupils of primary schools, Graz, etc.	4694		493 0.
"	"	"	Pupils of Grammar schools, etc., Graz.	569		376 33.3
"	"	"	Pupils of Keatsch. & Gymnasie, Graz.	1225	
83	"	Reich	Military Gymnasium pupils, Tiflis	246	
84	1882	Fox	American Indian pupils, Carlisle	8-22	250		237 94.8	6 2.4	4 1.6	?
85	"	Paulsen	Pupils of navigation school	76	152	119 78.35	19 12.5	14 9.25	?

1 The percentages foot up 102 per cent.

No.	Date.	Examiner.	Character of examined.	Age.	Persons. Eyes.	Em. per ct.	M. per ct.	H. m. per ct.	H. per ct.	As. and Amb.
86	1882	Borthen	Pupils of various schools, Thronbjerg	6-19	550	247=44.9	96=17.5	206=36.4		
87	"	Schubert ¹	Pupils of common school, Nuremberg	1012	626=34.2	505=27.6	699=38.2	?	Amb. 74
88	"	Kohert ¹	Pupils of var. schools, B. nos. Ayres.	6163	290 4.22		
89	"	Erlmann ²	School children (1870), St. Petersburg.	350	54 [±]	15 [±]		
90	"	Mittendorf & Derby, R. H.	Primary pupils, New York	203	[162 80.]	6 3	35 17.		
"	"	"	Grammar pupils, New York	698	[555 79.5]	[59] 8.5	[84] 12.		
"	"	"	Grammar pupils, New York	896	119 13.5		
"	"	"	College students, New York	201	69 35	[30] 10.		
91	1883	Nordenson ²	Pupils of Alsatian school, Paris	7-20	225	(319 70.6)	70 15.5	63 13.9	As. 411
92	"	Mauz	{ Gymnasium pupils, Freiburg	487	141 29.		
"	"	"	{ Grammar pupils, Freiburg	889	136 14.2		
"	"	"	{ Pupils of various schools, Freiburg	3982	452 11.		
93	"	Berlin and Rembold	Orphanage scholars, Stuttgart	7-15	282	103 18.		
"	"	"	Real-Gymnasium pupils, Stuttgart	41.7		
"	"	"	Gymnasium pupils, Suttgart	45.7		
94	"	Gaertner	Theological graduates, Tübingen	4-19	575	283 24.6	865 75.2	2 0.17		
95	"	Derby, Hasket	College graduates, Amherst	21-26	254	87 34.3	120 47.2	47 18.5		
96	"	Dobrowsky	High school pupils, Ural	212	[26] 12.0		
97	"	Dürr	Lyceum pupils, Hanover	[9-19]	345	46 13.3(?)	121 35.(?)	107 33.6	(Homatropia) (192 60.4)	
"	"	"	Seminary students, Hanover	[16-20]	96	9 9.37	82 33.3	36 37.5	(Homatropia) (55 57.3)	
98	"	"	Real-Gymnasium pupils, Hanover	[9-19]	271	81 30.4		
99	"	Hansen	Country school pupils, Kiel	10-14	808	26 1.58	51 3.4		Amb. 6
100	"	Reich	Girls' institute pupils, Tiflis	173	40 23.1	57 33.	63 36.4		As. 13
"	"	"	Infantry school pupils, Tiflis	292	52 17.8		
"	"	"	Military gymnasium pupils, Tiflis	252	61 24.2	49 19.4	129 51.2		As. 10
101	"	Motals	Type-setters, Paris	97	10 10.3	31 31.9	[17] 2.83	21 21.6	As. 30
102	"	Schtschepotjef	Middle school pupils, Astrachan	600	[482] 80.33	[99] 16.5	240 82.2	As. [3]
103	"	Schadow	Primary scholars, Island of Borkum	6-14	146	44 15.1	2 0.7		
104	"	Hadlow	Naval training sch'l pupils, Greenwich	15	1074	60 5.8(?)		
105	"	Tscherning	Recruits (educated), Copenhagen, etc.	2336	(+2 -2) (1825 78.12)	(420 18.)	(> +2) (91 3.9)	
"	"	"	Recruits (uneducated), Copenhagen	5187	(+2 -2) (4770 91.96)	(207 4.)	(> +2) (210 4.05)	
106	"	v. Reuss	Railway employes, clerks, etc, Vienna	273	150 54.94	86 31.50	37 13.55		
"	"	"	" " engineers, etc, Vienna	280	188 67.14	11 3.92	81 28.93		
"	"	"	" " trainmen, Vienna	421	307 72.92	25 5.93	89 21.13		
"	"	"	" " stationmen, Vienna	554	376 67.87	30 5.41	148 26.71		
107	1884	Schleich	Infants (newborn), Tübingen	2h.-8d.	350	0. 0. 0.	0 0 0.	(Atropine) 300 100.	

¹ Received too late for inclusion in summaries. ² Re-examination, vide 14. ³ Corneal astigmatism was measured with the ophthalmometer of Javal & Schiötz.

No.	Date.	Examiner.	Character of examined.	Age.	Persons. Eyes.	Em. per ct.	M. per ct.	H. m. per ct.	H. per ct.	As. and Amb.
108	1884	Schäfer	Deaf-mute pupils, Heidelberg	9-18	95	21 = 22.1	7 = 7.3	41 = 43.1	62 = 65.2	
109	"	Seggel	Recruits (common soldiers), Munich	20-24	1235	1423 56.7	347 11.4	1239 40.6	41
110	"	Ulrich	Recruits (volunteers), Munich	20-24	284	128 22.5	341 60.	71 12.5	(Atropine)	28
111	"	Bjerrum	Infants (newborn), Königsberg	4h.-14d.	102	0 0.	0 0.	204 100	
112	"	Besselin [Becker]	Infants (newborn)	7h.-14d.	87	23 26.5	8 8.4	(Atropine)	
113	"	Katelman	Girls' school pupils, Heidelberg	8-18	250	348 69.6	61 12.2	(Atropine)	As. 16
114	"	Seggel	Girls' school pupils, Heidelberg	5-18	869	485 65.8	62 8.4	?	(Atropine)	
115	"	Horstmann.	Soldiers (militia), Munich	15-35	17	9 26.5	0 0.	25 73.5	(Atropine)	As. 40
116	"	"	Military cadets, Munich	1118	(Atropine)	
117	"	"	Gymnasium pupils, Munich	350	(Atropine)	
118	"	"	Girls, boarding school pupils, Munich	298	(Atropine)	
119	"	"	Infants (newborn), [Berlin]	7-21d.	174	(Atropine)	
120	"	"	Young children, [Berlin]	1-2y.	50	10 10.	6 2.	(Atropine)	
121	"	"	Young children, [Berlin]	4-5y.	50	10 10.	6 6.	(Atropine)	
122	"	"	Indian girls, pupils, [Carlsruhe]	17-33	60	15 13.	13 13.	(Atropine)	
123	"	"	University students, Leyden	110	36 32.7	13 11.8	(Atropine)	
124	"	"	Municipal scholars, Lucna	6-16	470	527 56.23	291 31.	114 12.15	6
125	"	"	Boys, elementary scholars, Rome	5-10	266	532	142 26.7	7 1.2	Mix. As. 5
126	"	"	Boys, municipal scholars, Rome	6-15	76	152 (73)	6 4.	5 0.	
127	"	"	Girls, municipal scholars, Rome	6-13	350	700 (515)	91 13.	10 1.4	
128	"	"	Boys, municipal scholars, Sienna	7-15	220	440 (295)	67 15.2	30 7.	
129	"	"	Boys, mun. scholars, Castelflorentino	7-13	352	704 (422)	81 11.5	2 0.3	
130	"	"	Boys, elementary scholars, Cortialdo	7-14	102	204 (168)	16 7.8	2 1.	
131	"	"	Girls, element. etc., Naples, etc.	3-30	148	296 (221)	19 6.4	0 0.	
132	"	"	Boys, Lyceum, etc., Naples, etc.	6-22	722	1444 (1133)	190 13.1	14 1.	
133	"	"	Boys, Lyceum, etc., Naples, etc.	7-19	890	2074 (1752)	228 10.9	7 0.3	
134	"	"	Girls, elementary scholars, Trapani	8-15	415	156 31.2	131 15.8	1 0.1	
135	"	"	Girls, elementary scholars, Trapani	6-17	389	226 (175)	57 15.4	2 0.6	
136	"	"	Lyceum pupils, Strassburg	9-20	517	775 (575)	95 12.2	0 0.	
137	"	"	Upper Gymnasium pupils, Tübingen	14-18	60	246 47.4	190 36.6	(S1 15.6)†	
138	"	"	Medical students, Philadelphia	19-34	120	16 13.3	86 71.6	(S1 11.5)	
139	"	"	Medical students, Philadelphia	19-34	90	32 18.8	17 9.44	(87 26.)	131 72.77	

† In this group of Italian investigations the optometer of Moyné seems to have been used, and note made of only the absolute H. if any.

‡ The H. was measured with the ophthalmoscope, whether it was sought with this means among the apparent Em. does not appear.

Let us now see how far the methods above sketched have been applied in the investigations which have been made; what results, positive and negative, have been obtained; and how far we can formulate corrections for such as are incomplete, and by their application bring also the inferential results to the elucidation of our subject:¹—

Beginning our study of the data given in the preceding tables by a summing up of all that can be employed on the subject of the frequency of *Myopia*, we find that we have available records of 115 investigations or groups of investigations:—

Of 146,522 examinations (individuals or eyes) M. 28,483 = 19.44 p. c.

Separating, now, those investigations giving us data as to all conditions of refraction, we have as the result of 88 investigations or groups of studies:—

90,886 exam. : Em. 43,174 = 47.73 p. c., M. 19,750 = 21.87 p. c., H. 25,635 = 28.46 p. c.

In not a few of these cases the M. designates only *apparent myopia*, the Em. comprises all with normal vision, or at best those without *manifest* ametropia;² while under H. are included, for the most part, only the *H. manifest*—the cases where the *H. total* was sought being about counterbalanced by those in which only the *H. absolute* is given. Turning now to that series of studies in which *H. total* was sought and noted with care sufficient to obtain some approximation to it, and to exclude from M. the cases of spasm of accommodation simulating that anomaly, we gain the following results from 27 studies:—

16,160 examinations : Em. 3297 = 20.40 p. c., M. 2918 = 18.01 p. c., H. 9717 = 60.13 p. c.

This is excluding Erismann's 1889 = 43.3 per cent. H. m., Emmert's 3308 = 77 per cent. H. m., Pflüger's 800 = 44 per cent. H. m., Seggel's 1239 = 40.6 per cent. H. m., and many others with even higher percentages of H. m. It is also exclusive of Just's work, since the 1660 = 67. per cent. *non-myopic* eyes, although "almost all" hypermetropic are not with sufficient clearness stated to have been H.; and Weber's, Tscherning's, and Beselin's work, although doubtless accurately done, is excluded for lack of full data as to the results obtained.

¹ Lack of space excludes the detailed accounts of the methods and results of the individual investigations which were prepared to form the body of this paper: so also as to the tables of curves which would in some measure have supplied their place as to class-results, etc. The school totals are given in the tabulation which follows. Under Em. are included all not proven ametropic; under M. all whose apparent myopia was not shown to be fictitious; under H. m. all cases of hypermetropia manifest, whether absolute, facultative, or relative; and under H. all, where any attempt seems to have been made to discover the total hypermetropia, as well as those where the methods employed are not given. Astigmatic eyes are included under H. and M. respectively, when recorded as H. As. or M. As.; under As. when not so specified.

² Gardner found among 1082 pupils of Springfield, Mass., 81 = 7.5 p. c. M., 291 = 26.9 p. c. H. m. and 69 = 6.35 As. Report of School Committee, 1884.

Combining the cases where we obtain data as to the relations of *normal vision, apparent Em.* (*i. e.*, absence of H. m.), and Em. probably such, we find:—

5347 examinations: V=1. 4362=81.58 p. c., App. Em. 2578=48.21 p. c., Em. 147=27.51 p. c.

although among the Em. here noted, are still many not adequately tested, if at all, for latent H.

A similar comparison as to H. abs., H. m., and H. (generally ophthalmoscopically determined) gives:—

1378 examinations: H. abs. 59=4.29 p. c., H. m. 784=56.9 p. c., H. 1031=74.8 p. c. while in—

9344 examinations: App. Em. 4383=47. p. c., Em. 1742=18.6 p. c., H. m. 2076=22.2 p. c., H. 5284=56.5 p. c.

if we include Just's work, as well as others more definite. In 6873 examinations where App. M. constituted 1911=27.8 per cent., these figures were reduced to 1415=20.6 per cent. by measures calculated to eliminate more or less completely the accommodative spasm. Similar comparisons could be made on many points; but this is sufficient to show the direction and extent of the corrections which would have to be applied to many of the studies before us in order to approximate accurate results. The actual application of such corrections can hardly be profitably made, except to very extensive summaries, and even then with great reserve; yet we will probably be fully justified in claiming that, among the whole number thus far examined, more than 145,000 in all, not more than 20,000, or 14 per cent., were actually myopic. Similarly, of about 90,000 examined as to the relative frequency of the various conditions of refraction, some 22,000 were approximately Em. (25 per cent.), 13,000 M. (14 per cent.), 30,000 H. m. (33 per cent.), and 55,000 H. (60 per cent.).¹

Leaving this field of conjecture, and turning to the data showing the refraction at various ages, we will first study the question of the refraction of the new-born infant. Eight investigations are on record:—

	Infants.	Eyes.	Em.	M.	H.
1861. v. Jaeger,	(100+)	100	5 = 5 p. c.	78 = 78 p. c.	17 = 17 p. c.
1880. Ely,	111	154	21 14 "	27 18 "	106 69 "
" Horstmann,	20	40	8 20 "	4 10 "	28 70 "
1881. Königstein,	281	562	10 2 "	0 0 "	552 98.2 "
1884. Schleich,	150	300	0 0 "	0 0 "	300 100 "
" Ulrich,	102	204	0 0 "	0 0 "	204 100 "
" Bjerrum,	87	(87)	23 26.4 "	3 3.4 "	61 70.1 "
" Horstmann,	50	100	10 10 "	2 2 "	88 88 "
	901+	1547	77 = 5 p. c.	114 = 7.3 p. c.	1356 = 87.6 p. c.

All authorities are agreed that *full paralysis of the accommodation* by a mydriatic is an absolute essential in this difficult field of investigation; and we must accept Horstmann's statement that this is attainable with

¹ Study of the degree of refraction error, of anisometropia and of many other points of cardinal importance must be reserved for a later paper.

difficulty, even when employing a solution of atropiæ sulph. 1:100. Jaeger's work, done in the earliest days of the ophthalmoscope, probably lacked that exquisite skill in the use of the instrument which later distinguished this master of ophthalmoscopy; and, further, was done without employment—certainly without the complete employment—of a mydriatic. His parallel work in the study of the dimensions of 70 eyes of infants after death, showed no instance of axial myopia; and the later studies force us to decide against his assumption of an undeveloped suspensory ligament or of a usual forward displacement of the lens, causing a *temporary* myopia of the new-born—a view negatived also by the rest of his examinations of the eyes of older children. We must, therefore, exclude Jaeger's study (albeit with much hesitation) as incompetent evidence in the matter before us. The need of "full paralysis of accommodation" leaves room for questioning of the results obtained by weak mydriatic solutions, since Horstmann failed in at least 7 instances to secure even full dilatation of the pupil with 1 per cent. solution of atropine repeatedly used. Bjerrum's finding of 23 Em. and 3 M. after (a single?) employment of atropine (0.25 per cent. solution) cannot be considered conclusive; and Ely's results of 17 per cent. Em. and 11 per cent. M. among 105 eyes under the influence of a solution (gr. ij ad f̄3j) of atropine, and 8.2 per cent. Em. and 32.6 per cent. M. among 49 eyes under a weak atropine solution (gr. j ad Oj) must be similarly viewed. Accepting them conditionally, however, and doubling Bjerrum's figures to obtain the number of eyes studied by him (a procedure apparently justifiable), we gain as the result of these studies, exclusive of v. Jaeger's:—

Of 1534 eyes of infants, Em. 95 = 6.19 p. c., M. 39 = 2.54 p. c., H. 1400 = 91.26 p. c.

Only the 6 myopic eyes of Horstmann's two studies can be considered as well-established cases of M.; yet the success of Königstein and Ulrich with atropine (0.1 per cent.), and of Schleich with 0.5 per cent., in doing away with all apparent M. in their studies, lends confirmation to other results where weak mydriatic solutions were employed.

Passing on to the consideration of the refraction of children more developed, but not yet of school-age, we find, exclusive of v. Jaeger's work, only four investigations.

	Eyes.	Em.	p. c.	M.	p. c.	H.	p. c.
Koppe, Kindergarten pupils, 30	60	1	1.7	0	0	59	98.3
Horstmann, children:							
(8-24 months), 57	96	13	13.5	9	9.4	74	77
(1-2 years), 50	100	10	10.10	6	6	84	84
(4-5 years), 50	100	13	13	13	13	74	74
Total children (8 m.-5 yrs.), 187	356	37	10.4	28	7.86	291	81.75

¹ Koppe's results in the Kindergarten are impeached by himself, since he may have been measuring his own hypermetropia and not that of the observed eye. All but 18

Proceeding now to the other end of the scale of life, we find the one study of Cohn on the eyes of aged persons, mountaineer villagers in Silesia :—

100 persons (60–84 years), 200 eyes, Em. 49 = 24.5 p. c., M. 25 = 12.5 p. c., H. (m.)
114 = 57 p. c. ? 12 = 6 p. c.

A study of much interest, but of insufficient extent to be conclusive, while the individuals tested were within the region of the “hypermetropia aquisita,” and of myopia due to the cataractous swelling of the lens, the “second sight” of common parlance. The other investigations of the eyes of adults have generally more bearing upon the question of the ophthalmic hygiene of various handicrafts than upon our subject, until we reach the investigations of the eyes of recruits, almost invariably young men from 20 to 24 years of age. Considering only the common soldier, whose eyes have been subjected to primary school-work alone, examinations as to visual acuity have been extensive and interesting. Thus Herzenstein found among 27,682 men of various arms, binocular vision above normal with Junge’s test-types in from 71.6 per cent. to 76.6 per cent. of various garrisons about Orel and Cracow ; while Rumschewitsch among 9882 recruits in the Kiew district, found 70 per cent. with V. = $\frac{120}{XL}$ to $\frac{60}{XL}$,

26.4 per cent. with $\frac{60}{XL}$ to $\frac{40}{XL}$, 4 per cent. with V. = $\frac{20}{XL}$ to $\frac{40}{XL}$, and 0.6

with V. $\frac{1}{XL}$ to $\frac{20}{XL}$.¹ The examinations bearing more strictly upon the refraction are numerous, but rarely of both convincing accuracy and fulness.² Seggel’s study of the common soldiers of the Munich garrison is about the only one fully competent in its testimony, and from its extent it can well stand alone.

1526 common soldiers, 3052 eyes, Em. 1425 = 46.7 p. c., M. 347 = 11.4 p. c., H. m.
1239 = 40.6 p. c., Amb. 41.

Allowance must, of course, be made for the cases with latent H. among those noted as Em. in order to approximate full accuracy ; a correction which cannot, however, be accurately formulated.

eyes, however, appeared H. > 1. D., the amount which he estimates that he manifested in this investigation, and only 2, including the 1 Em., were H < 1. D.

The Kindergarten and El. pupils were examined by the ophthalmoscope alone. In the other schools he obtained (counting As. with M. and H.) :—

by glasses,	989 pupils,	Em. 219 = 22.1 p. c.	M. 269 = 27.8	H. m. 501 = 50.7 p. c.
by ophthal.	1059 “ “	188 = 17.7 “	M. 240 = 22.6	H. 606 = 57.2 “
by oph. and gl.,	1059 “ “	112 = 10.6 “	M. 220 = 20.8	H. 711 = 67.1 “

¹ The question as to the illumination under which these results were obtained is all-important ; yet many investigators are strangely silent in this regard. In sunlight $\frac{120}{XL}$ would indicate no sharper vision than $\frac{20}{XV}$ in a well-lighted room.

² Ljubinsky is cited (Nagel’s Jahresbrt. xiv. p. 195) as having found, among 7312 sailors, Em. 48.7 p. c., M. 28 p. c., H. 43.1 p. c., and As. 2.2 p. c. (120. p. c.).

Of the data that remain all are complicated by the question of "school myopia," into which it is impossible here to enter. The persons examined were pupils of schools of every grade, generally fully engaged with their school-work, and showing at once the temporary and the permanent marks of its effect upon their eyes. Yet much information can be obtained from their study, and we will proceed to a rapid review of the refraction in the schools of various grades:

Summing up, as before, the results of the twenty-six investigations as to myopia, we learn that in 23,315 examinations 1582 = 6.79 per cent. M. were found. A discrimination between the real and the apparent M. in these results would lower somewhat the M. percentage, although the figures of the first two studies on the list contain no myopes below 1. D. in grade, and the last study, only those greater than 1. D.

Passing to the other conditions of refraction we find—

In 22 investigations	13,929 examinations :	Em. 857 = 63.59 p. c.,	M. 1582 = 11.36 p. c.
		H. 3764 = 27.04 p. c.	
" 10 "	3,358 "	Em. 491 = 14.62 p. c.,	M. 137 = 4.08 p. c.
		H. 2564 = 76.06 p. c.	

In the second series are placed those studies in which the H. t. was sought, although in Beselin's work only the H. > 1. D. is reported, and here are also included Just's results, although not entirely definite in their statement—any error in the one being probably neutralized by the other. From five investigations, exclusive of Just's, we obtain:—

1835 examinations :	App. Em. 767 = 41.79 p. c.,	Em. 371 = 20.22 p. c.,	App. M. 126 = 6.86 p. c.,	M. 58 = 3.16 p. c.,	H. m. 970 = 52.86 p. c.,	H. 1368 = 74.55 p. c.
---------------------	-----------------------------	------------------------	---------------------------	---------------------	--------------------------	-----------------------

These figures comprehend the results of Cohn's study in the mountain village of Schreiberhau, in Silesia, when atropine (in substance) was employed in most of the eyes. As various critics take diverse views of the question of the refraction under mydriatics, a few remarks upon the subject seem in place. Donders's definition of refraction fittingly opens the subject: "By *refraction of the eye*, we understand its refraction in a state of rest; that is, the refraction which the eye possesses in virtue of its form and of that of its component parts, independently of muscular action, independently of accommodation. The term, therefore, applies to the refraction of the eye whose muscles of accommodation are inactive or paralyzed (*for example under the influence of atropia*), to the refraction also of the dead but as yet otherwise unaltered eye."¹ In discussing emmetropia, however, Donders assumes a "*tone of accommodation*" which can be done away only by a mydriatic, and concludes with the often-quoted expression: "Consequently, the actually emmetropic vision requires in a certain sense a minimum of H., and that minimum is capable of no accurate taxation, because to the tone itself a certain latitude, perhaps from $\frac{1}{100}$ to $\frac{1}{40}$, must be allowed." Whether his view on this point remains unchanged we do

¹ Donders, Accommodation and Refraction of the Eye, p. 80. (Italics ours.)

Elementary Schools, etc.

Examiner.	Examined.	Age.	Pupils.	Eyes.	Em. per ct.	M. per ct.	Hm. per ct.	H. per ct.	
Cohn . . .	Country Primary . . .	6-13	1486	1408=94.7	22= 1.4	16= 1.1	M. > 1. D.
Cohn . . .	City Elementary . . .	6-14	4978	4245 85.3	334 6.7	127 2.5	M. > 1. D.
Erismann . . .	City Preparatory	59	11 18.6	8 13.6	40 67.8	Atropine.
Cohn . . .	Village Primary . . .	6-13	(240)	480	14 3.	4 0.9	370 77.	438=91.2	
v. Hoffmann . . .	City Preparatory . . .	6½-9½	249	147 59.	4 1.6	13 5.2	88 31.3	
Callan . . .	Negro Primary . . .	6-13	270	18 6.6	2 0.84	213 78.9	250 92.	
Conrad . . .	City Preparatory . . .	[6-9]	568	155 27.3	28 5.	101 17.6	383 67.4	
Spalding . . .	City Primary . . .	6-10	1200(+)	84 7.	
Pflüger . . .	City Primary . . .	7-10	808	383 84.9	27 3.35	409 50.6	
Emmert . . .	Elementary . . .	6-12	367	734	103 14.	32 4.86	596 81.2	
Loring & Derby . . .	City Primary . . .	6-12	410	348 84.9	28 6.8	34 8.3	
Koppe . . .	City Elementary	(103)	206	53 25.7	5 2.4	126 61.1	
Koppe . . .	City Preparatory	(136)	273	37 14.	20 7.5	173 70.	209 78.5	
Haenel . . .	City Primary (classes)	682	1364	784 57.47	226 16.68	348 25.49	
Just . . .	City Primary	(281)	563	(433 77.)	59 10.5	59 10.5	[492 87.5]	
Smith . . .	City Board School . . .	7-13	1636	85 5.3	
Netolitzka . . .	City and Village Primary	4934	493 10.	
Dennett . . .	Village Primary . . .	4-10	530	430 81.1	21 3.	58 10.9	
Florschütz . . .	City Primary, 1873 . . .	6-15	1476	1053 71.3	198 13.4	60 4.1	
Florschütz . . .	City Primary, 1877 . . .	[6-15]	1616	90 5.57	
v. Reuss . . .	City Primary . . .	6-11	257	129 40.4	29 11.2	99 37.1	
Risley . . .	City Primary . . .	5-12	(164)	328	23 7.01	13 4.	289 88.1	
Borthen . . .	Common . . .	7-13	133	64 48.1	8 6.	61 45.8	
Mittendorf & Derby . . .	City Primary	203	[102 80.]	6 3.	85 17.	
Nordenson . . .	City Primary . . .	7-13	79	61 77.2	5 6.3	13 15.2	
Schadow . . .	Country Primary . . .	6-14	(146)	192	44 15.1	2 0.7	240 82.	
Beselin . . .	City Primary . . .	5-9	131	70 53.4	4	49 37.4	{ Am. > 1, { Em. = 1, 1, 1.

not know; but can hardly imagine so careful an observer to be still in error on this simple point. The question before us is whether this common residual tension is an *organic tonicity* or an *habitual action*. That no such "tone" is ascribed to the myopic eye is curious, if the first supposition holds good. Is there, then, in the eye, a tension of the accommodation which *cannot* be voluntarily relaxed? It may be confidently asserted that an emphatic negative will be given by *every observer who has carefully tested the matter*.

The assumption of such a "tone," like the hypothesis of *negative accommodation*—really another phase of the same question—lacks the first *positive* proof in its support. Negative evidence alone, cases of *habitual spasm* of the accommodation, can be adduced in its favor; and a single positive fact, did it stand alone, would scatter such support to the winds. The investigation before us fully answers the question, although its testimony has been generally overlooked. Cohn states: "*In 34 cases with H. m. in degree from $\frac{1}{8}$ to $\frac{1}{2}$ there was no slightest increase of the H. after atropinization.*" It may be remarked that *full paralysis* of accommodation was *proven* to have been attained in these cases, although in four others who remained apparently emmetropic, Acc. = $\frac{1}{15}$, $\frac{1}{24}$, $\frac{1}{30}$, and $\frac{1}{35}$, respectively, was found remaining. Similar proof, were any needed, would be found in the investigations of Roosa (53) and Dürr (98). How two opinions on the subject can exist is the only matter of surprise to the writer, who has in hundreds of cases given the *full correcting-glass* as determined under a mydriatic pushed to complete paralysis of accommodation (and seen it done in thousands), and has rarely seen it necessary to reduce the strength of the glass because of persisting recurrence of the habitual spasm.¹ His own low H., twice measured under full mydriatic paralysis, is entirely relaxed at will, and always taken into account in ophthalmoscopic measurements. The only noteworthy point, therefore, as to the admission of Cohn's atropine work in our present study, is that *especial weight* is to be given to its showings, since in it was employed a means of accuracy lacking in others.

The group of schools in which were found the results which have been given, practically represent the first three years of school-life; and it is from this point that the course of the German Gymnasium takes its start. Intermediate schools exist in Germany as well as elsewhere, but they differ as to the upper limit of their curriculum, and in the aims and the class of the pupils attending them, rather than in the age of their scholars. While the distinction between these and the gymnasia is, therefore, very important in its bearing on the question of school-myopia, to our investi-

¹ The necessity, or, indeed, the advisability, of carrying this matter in practice to its extreme logical conclusion is a point which most ophthalmologists are by no means prepared to admit. Its *practicability* is the only question here raised, and is, in the writer's opinion, too fully demonstrated to need more extended proof.

Examiner.	Schools.	Age.	Per. Eyes. 800s.	Em. p. c.	M. p. c.	VIII.	VII.	VI.	V.	IV.	III.	II.	I.	Hm. p. c.	H. p. c.
Cohn.....	Realschule.	10-22	1141	816=75.9	225=19.7	16= 9.	44=16.7	51=19.2	50=25.1	41=26.4	23=44.	44=1.9	
Cohn.....	Gymnasia.	9-22	1195	896 72.5	314 26.2	31 12.5	41 18.2	52 23.7	73 31.	61 41.3	54 55.8	9= 2.9	
Thillius.....	Gymnasium.	314	175 55.7	96 30.6	10 5.4	16=26.2	33 33.	20 44.4	45 50.6	32 54.2		
Schultz.....	Various.	431	169 36.9	1=14.3	16=26.2	10 5.4	22 36.7	13 38.4	20 44.4	45 50.6	32 54.2	1825 43.	
Erisman.....	Gymnasium.	4239	1110 25.6	1284 29.7	124 15.8	197 25.4	262 30.7	228 32.4	206 41.3	127 42.	110 43.8	25 7.	
Cohn.....	Various.	6-22	361	187 52.	125 35.	5 13.	13 32.2	14 27.7	21 56.	33 62.4	27 65.	12 42.	
Kruger.....	Gymnasium.	94-19	203 466	150 39.	162 40.	6 7.	13 32.2	21 28.4	22 32.	25 32.1	53 63.3	22 62.5	83=20.6	
v. Hoffmann.	Gymnasium.	94-19	256 512	140 27.3	187 36.5	16 19.	28 23.7	21 28.4	22 32.	25 32.1	53 63.3	22 62.5	[101 19.7]	171 33.8
v. Rens.....	Gymnasium.	409 818	243 29.7	375 45.8	41.	48.	48.	40.	50.	61.	68.	158 19.3	[184]
v. Rens.....	Gymnasium.	389	144 37.	106 42.7	37.	42.	46.	45.	55.	69.	75.	69 17.7	
Ott & Ritze- mann.....	Gymnasium.	122 244	124 61.	96 39.3	[15] 24.8	[25] 40.3	[16] 30.	[18] 47.	[18] 47.	[14] 54.	[7] 68.	24 10.	?
Ott.....	Realschule.	12-14?	164 324	146 45.	97 31.1	14 5.5	[15] 14.9	4 12.9	122 37.7	?	
Dor.....	Realschule.	10-19	430	279 64.9	125 29.	13 17.1	19 28.7	19 26.4	23 54.9	23 41.1	12 40.	11 68.	18 4.18	
Conrad.....	Gymnasium.	8-20	1234 2468	754 30.65	650 26.34	36 8.7	54 14.3	81 18.6	200 34.	155 37.4	124 51.7	256 10.37	1058 42.9
Spading.....	Grammar.	10-13	1000(+)	439 43.9	[100] 16.1	404 40.4	
Flüger.....	Various.	11-23	1065	131 15.1	14 6.3	12 5.8	20 10.	22 14.4	26 26.6	23 34.5	34 67.	2712 76.1	
Emmert.....	Various.	1781 3562	319 8.95	510 14.3	
Loring & Derby.....	Various.	1855	1183 63.8	420 22.6	252 13.6	?
Scheidt.....	Latin classes	924 448	134 29.9	155 34.6	21 21.8	30 25.8	44 41.5	46 42.6	14 63.6	28 6.25	
Scheidt.....	Gymn.	199 298	53 17.8	216 72.5	56 59.6	57 75.	50 80.6	63 80.3	29 9.7	
Koppe.....	Gymn.	396 792	75 9.7	200 26.	18	14	31	41	33	31	328 44.2	502 64.5
Kotemann.....	Realschule.	8-21	413	240 58.	157 38.	32	45.3	40.2	48.2	61.2	10 2.4	
Kotemann.....	Realschule.	232	[90] 26.	16 7.	14 7.	21 45.	29 45.	45.	45.	
Classen.....	Realschule.	492	[66] 41.29	29.	24.	40.	46.	71.	50.	
Backer.....	Gymnasium.	287	154 53.	100 35.	4.	100.	>1. D?
Agnew.....	College.	549 1097	651 59.	333 30.3	179 26.7	71 35.	35 30.7	30 60.	18 37.5	105 9.6	?
[Cheatham] & Mathewson.....	Techn. Inst.	300 600	372 62.	79 12.5	126 21.	?
Derby, Has- kett.....	Coll. Freshm.	122	145 45.2	36 29.5	67 20.9	?
Derby.....	Coll. Freshm.	15-24	321 650	114 35.9	
Niemann.....	Gymnasium.	325 650	[318] 49.	23.	29.	39.	63.	58.	75.	
Niemann.....	Paedagogium	388 776	[340] 44.	25.	27.	42.	42.	56.	70.	
Ott.....	Lyceum.	7-21	2032	1300 64.	568 28.	22 10.3	51 19.6	69 21.3	185 42.6	175 39.6	63 18.1	87 4.2	?
Ott.....	Gymnasium.	120 240	90 37.5	98 40.8	52 21.6	
Hael.....	Gymnasium.	177 354	[145 40.9]	209 59.8	
Hael.....	Gymnasium.	9-21	480 960	450 46.9	449 46.8	44 31.	56 33.7	83 47.9	75 48.1	46 54.8	89 62.2	53 68.8	54 5.62	?
Hael.....	Kreuzschule.	8-21	554 1108	460 41.5	564 50.	24 29.6	47 33.1	73 47.4	78 51.3	72 46.7	162 61.5	108 60.7	86 7.76	
Hael.....	Realschule.	8-17	209 418	204 48.5	161 38.5	8 25.	23 25.8	73 56.1	28 31.1	13 26.	16 61.5	52 12.4	
Hael.....	Seminary.	13-21	218 436	164 37.5	23 43.9	49 47.6	42 50.6	42 50.6	31 59.6	32 64.	25 56.8	32 7.6	

Examiner.	Schools.	Age.	Per. sons.	Em. p. c.	M. p. c.	VIII.	VII.	VI.	V.	IV.	III.	II.	I.	Hm. p. c.	H. p. c.	
Haezel	Seminary	13-20	131	292	103=39.3	134=51.1	23=40.3	24=54.5	16=40.	21=80.8	28=77.7	24=9.2		
Haezel	Secondary	8-16	1035	2060	1028 49.9	612 29.7	76=25.2	139=24.4	109 22.8	156 36.2	420 20.4		
Just	Gymnasium	194	388	167 40.4	195 50.3	26 37.1	33 32.8	48 52.	38 76.	39 67.2	27 7.	[103=49.7]	
Just	Realschule	293	586	264 45.	266 45.4	23 28.	31 29.2	46 52.3	87 58.	44 62.	24 4.1	[266 54.6]	
Koelmann	Gymnasium	8-17	253	566	182 31.6	110 19.4	12 9.5	23 18.2	27 22.1	32 22.2	14 39.	2 25.	273 48.8	
Knibbe	Normal Sch.	18-23	337	72 20.	?
Dannert	Various	10-19	603	424	70.8	74 12.8	23 9.8	19 10.5	18 16.5	10 19.6	3 33.3	76 12.6	
Emmet	Gymnasium	6-18	203	406	36 8.9	52 12.8	49.	318 78.6	
Florschütz	Gymnasium	9-20	177	407	91 51.4	24.2	37.	69.7	12 89.7	7 77.	6 3.4	
Florschütz	Realschule	10-18	260	122	47.	110 42.5	33.	33.3	34.5	37.5	55.5	19 65.5	5 1.9	
Florschütz	"	1871	280
Florschütz	Gymn. 1871	182	101 34.8
Florschütz	Gymn. 1871	44	90 48.4
Florschütz	Seminary	17-20	472	31.2	66 14.6	161 34.1
Wolter	Mil. Gymn.	609	1018	355 58.4	403 39.67	72 7.07	[105 +]
Wolter	Realschule	384	2094	250 33.3	240 31.67	50 7.07	[70 +]
Wesley	Various	9-21	149	298	283 11.4	319 16.2	1503 71.9
Natolicka	Gymnasium	119	298	182 44.6	149 50.2	11 32.	23 33.	38 51.	28 54.	22 85.	16 5.3
Natolicka	Realschule	254	136 33.5	149 50.2	15 29.8	7 16.3	8 21.	12 29.2	7 31.8	13 40.9	11 50.	35 13.8
Natolicka	" & Gymn.	1295	370 30.7 49	51	38	47	54	53	58	26	147 35.2
Barthen	" & Gymn.	417	183 43.9	88 21.1
Mittendorf	Grammar	698	[162 80.]	60 8.5	[84] 12.
Mittendorf	Grammar	896	119 13.5
Mittendorf	College	201	[111 55.]	69 35.
Nordenson	Lycium	10-20	147	667	97 667	32 21.8
Manz	Gymnasium	[487]	141 29.
Manz	Secondary	[280]	52 18.5	7 11.
Manz	Secondary	[603]	74 12.2	0 0.
Dobrowolsky	212	[23] 12.
Dürr	Gymnasium	[9-19]	318	25 7.86	101 31.7	9 15.5	12	8	11	22	33	23	107 33.6	Homat'p	
Dürr	Seminary	[16-20]	96	9 9.4	32 33.3	192 60.4	Homat'p	
Dürr	Real-Gymn.	[9-19]	271	81 30.4	2 12.5	9 18.	8 16.	15 38.5	23 33.9	22 50.	10	36 37.5	55 57.3	
Hansen	Girls' Inst.	10-14	868	1610	26 1.58	51 3.4	4 1.
Reich	Girls' Inst.	173	40 23.1	57 33.	1 17.
Reich	Mil. Gymn.	232	6.7
Reich	Mil. Gymn.	232
Becker	Grammar	8-18	250	500	348 69.6	61 12.2
Becker	Grammar	607
Baselin	Lycium	8-18	517
Hofmann	Lycium	9-20	246	47.4	190 36.6
Schleich	Upper Gymn.	14-18	60	120	16 13.3	86 71.6

>1. D.

75 15.

>1. D.

105 17.3

81 15.6

14 11.6

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

gation it is hardly more than incidental. We will group these institutions, then, by the side of the so-called "higher schools," which aim at liberal education; only premising that we are dealing now with pupils between the fourth and the twelfth school year, or between about the ninth and the eighteenth years of life.—

Combining the foregoing figures, we find as the result of examinations as to myopia in 50 gymnasiums and lyceums, 14 realschule, 7 seminaries, and about 50 other schools:—

In 55,342 examinations (individuals or eyes) 14,907 = 26.94 p. c. myopic.

Studying, now, those in which all conditions of refraction were sought, we obtain:—

In 42,763 exam. Em. 15,996 = 37.40 p. c., M. 11,757 = 27.49 p. c., H. 13,145 = 30.74 p. c., the H. probably here, as before, fairly representing the manifest hypermetropia, and the other figures the apparent myopia and apparent emmetropia, respectively. In the studies where the H. t. was sought, we find:—

In 9965 examinations Em. 2159 = 21.66 p. c., M. 2149 = 21.56 p. c., H. 5587 = 56. p. c.,

Here Just's results are again excluded, as well as those of Weber, of Becker, and of Beselin, where the figures given are confessedly incomplete and not available for our purpose; while Erismann's 1825 = 43 per cent. H. m., Pfüger's 404 = 40 per cent. H. m., Emmert's 2712 = 76.1 per cent. H. m., and Kotelmann's 273 = 48.8 per cent. H. m. are, of course, omitted. Studies of the relations of normal vision, apparent Em. and Em., of apparent M. and M., and of H. absolute, H. manifest and H. total, give results closely similar to those already set forth, and need not be here repeated in detail.

The data as to students of more advanced grade are limited, but doubtless worth summarizing:—

	Age.	Persons.	Eyes.	Em.	M.	H. m.
Cohn, Univ. of Breslau,	17-27	410		134=32.7 p. c.	244=59.5 p. c.	16=3.7 p. c.
Cohn, Medical students,		108	216	81 37.5 "	116 54. "	19 8.8 "
Gärtner, Theol. " Tübingen,	+19	713	1426	324 22.9 "	1096 76.9 "	2 0.2 "
Derby, H., Graduates, Amherst,	19-23	254		87 34.3 "	120 47.2 "	47 18.5 "
Collard, Univ. of Utrecht,	17-38	410	820	389 47.4 "	286 35. "	155 19.2 "
Van Anrooy, Univ. of Leyden,	17-38	470	939	527 66.23 "	291 31. "	114 12.15 "
Randall, Med. students, Phila.	19-34	71	142	51 48.8. "	54 10.5 "	37 26. "
	17-38	2436	4207	1593=37.86 p.c.	2207=52.46 p.c.	392=9.31 p. c.

In the first examination of this series H. abs. alone was noted; in the others the H. manifest was sought, but only in that of the writer, which ends the list, does the attempt seem to have been made to find the H. total by the ophthalmoscope or other means. In this last case the results were:—

71 Persons. 142 Eyes. Em. 30 = 21.1 p. c. M. 17 = 12.4 p. c. H. 93 = 65.5 p. c.

And for all—

90 Persons. 180 Eyes. Em. 32 = 18.8 p. c. M. 17 = 9.44 p. c. H. 131 = 72.77 p. c., figures in marked contrast to those above.

Looking backward over the long array of investigations which have been here grouped together, a few words of general comment seem permissible before passing to our conclusions. It is very evident that the majority of the studies cited can make no claim to being the unimpeachable foundation for our investigation of the refraction of the human eye; indeed few have been undertaken with the solution of any such question in view. The details of the methods employed in each examination would make clearer, perhaps, than is now the case, the true status and value of each; but lack of space excludes them, and the writer is unwillingly forced to group the studies and marshal their results as his study of the original data indicates is correct, and to set them forth apparently solely upon his own authority. Careful study of the original records will show, however, that the records have been in no instance classified, recast, or emended, arbitrarily; and if any judgment has been passed not fully upheld by the data furnished by each investigator, it has only been in holding as competent some of the investigations which are not *a priori* clearly self-condemned. In view of the great preponderance of the hypermetropic refraction in all investigations where its presence has been sought with a care in any degree adequate, it must be as evident to others as it is to the writer, that many of the studies giving low percentages in this matter cannot be correct. Such a conclusion, however, would by many be deemed a result of reasoning *a posteriori*, and has not been here drawn. For the figures given, unimpeachable accuracy cannot be claimed, especially when the sources from which they are derived are rarely free from palpable errors; and confusion and error are too easy of entrance to have been entirely excluded in the writer's own work. All that repeated careful revision and verification can do has been sedulously employed to secure correctness, and where possible the data have been submitted for verification to the investigators themselves, to many of whom the writer's thanks are therefore due.

It is then evident, that for the solution of the broader questions of the prevailing condition of refraction and other similar generalizations, few of the investigations thus far made are fully available; and it is to be hoped that future students of the subject will recognize these shortcomings and furnish us with fuller, broader, and more accurate data. The opportunities to employ mydriatics in extensive studies will continue to be very rare; it is therefore incumbent upon those who shall have such chances, if they would contribute fully to the elucidation of this important subject, to furnish at the same time data as complete as possible as to the vision without and with glasses, the accommodation and the apparent refraction, before the mydriatic, as well as the intraocular condition and the functional comfort of the eyes; and to furnish proof, where possible, that the mydriatic had been pushed to *full* paralysis of accommodation. Thus only can the strict requirements of the subject be met. For those

who are debarred from the use of mydriatics, combination of all other methods of study is necessary, if they desire to render as slight as possible the doubt which must always attach to determinations of the refraction made in eyes with unparalyzed accommodation.

The following conclusions seem fully upheld by the results of the investigations thus far made:—

1. Myopia is almost unknown in infancy and very infrequent before the beginning of school-life. In the earlier school-years its percentage is still low and it is only in the advanced classes, especially of the German schools, that it ever attains to a preponderance. It has been found in not more than $39 = 2.54$ per cent. of 1534 eyes of infants, in not more than $28 = 7.86$ per cent. of 356 eyes of children under the school-age, and in only $1582 = 6.79$ per cent. of 23,315 eyes of children examined during the first three school-years—figures which more accurate methods might have made lower. Among 3052 eyes of young men, upon whom the school influence had not been excessive, it was found in $347 = 11.4$ per cent.—a percentage which probably oversteps the maximum which it is likely to attain outside of the schools.

2. Hypermetropia is the enormously preponderating condition in infancy and early childhood, and the first years of school life witness little reduction in its proportion. Outside of the schools it remains by far the most frequent refraction throughout life, and in the schools it is decreased by the change of eyes to the myopic refraction in a degree apparently varying according to the circumstances calling into existence that defect. It was found in $1400 = 91.26$ per cent. of the 1534 eyes of infants examined, in $291 = 81.75$ per cent. of the 356 eyes of young children, and in $2564 = 76.$ per cent. of the 3358 eyes of children in the elementary school years, among whom it was sought with adequate care. So also in the higher schools, it constituted at least 56 per cent. of the whole number of eyes studied by competent methods, being found in 5587 of the 9965 examined.

3. Astigmatism has been rarely sought with care, and the data with regard to its frequency are not sufficiently wide to justify definite conclusions. The findings of the studies where it has been well looked for, concur with the clinical work in indicating a measurable degree of astigmatism (0.5 D. or more) in the majority of ametropic eyes.

4. Emmetropia in a mathematically strict sense has probably no existence. Approximate emmetropia (Am. $< \pm 0.5$) is infrequent in all ages, probably at no epoch exceeding 10 per cent. Its apparent proportion is swollen by the array of eyes "*not proven*" ametropic, and we have but few studies where the accommodation has been with certainty set aside and its existence fairly well shown. Cohn among 299 atropinized eyes proved in no single instance its presence. Under homatropine Hansen found it in but 26 of 1610 eyes, and Dürr in 30 of 414 eyes: it consti-

tuted at most $60 = 2.6$ per cent. of these 2323 eyes. Among the infants and young children $135 = 7.36$ per cent. of the 1834 examined under atropine may have been emmetropic; and Roosa's brief study indicates that it is probably as rare in adult life, even when perfect function apparently proves its presence.

5. The question of what is the *normal refraction* of the human eye is still an open one, and further material on the subject and closer study of the data in hand will be necessary before drawing conclusions as to it. Much light will be thrown upon the question by studies like Risley's of the relations of normal vision, intraocular health, and functional comfort, to the refraction. For the present the conclusion of this author, whose work stands almost alone, may be accepted. "The emmetropic is the model or standard eye—since emmetropia is shown not only to remain nearly constant in percentage throughout the school life, but that it is also the condition of health, and withal enjoys the highest acuity of vision and the greatest freedom from pain."¹ Yet hypermetropia is the prevailing—almost the exclusive—condition of the refraction among most animals, among children, among uncivilized peoples, and among all eyes uninjured by the educational process.

1806 CHESTNUT ST.

BIBLIOGRAPHY.

1. James Ware. Observations relative to the near and distant sight of different persons. Philosoph. Trans. Royal Society, London, 1813, 1. p. 31.
2. Szokalski. Ueber die hygienische und therapeutische Anwendung des Schutzbrillen Vierteljahrschrift f. d. prakt. Heilkunde, Prag, 1848, p. 139.
3. J. H. Schürmayer. Handbuch der medicinischen Policei, 2d ed. Erlangen, 1856.
4. Ed. v. Jaegar. Ueber die Einstellungen des dioptrischen Apparates, Vienna, 1861.
5. Ruete. Untersuchungen über die Augenkrankheiten bei Schulkindern. Zeitschrift, f. Med. Chir. u. Geburtsh., v. p. 223, Leipsic, 1866.
6. Reck.
7. Alexander. } Untersuchungen der Augen von 10,060 Schulkindern, Leipsic,
8. Gaertner. } 1867.
9. Cohn.
10. Cohn. Die Augen der Breslauer Studenten, Berlin. Klin. Wochenschr., No. 50, 1867.
11. Cohn. Die Augen der Breslauer Schriftsetzer, Berlin. Klin. Wochenschr., No. 50, 1868.
12. Hugo Thilenius. Untersuchung der Kurzsichtigkeits-Verhältnisse auf dem Rostocker-Gymnasium. Inaug. Diss. Rostock, 1868.
13. Schultz. Redogörelse för Elementarläroverken i Upsala, etc., p. 29, Upsala, 1870,
14. F. Erismann. Ein Beitrag zur Entwicklungs-Geschichte der Myopie, Arch. f. Ophth., xvii. p. 1, 1871.
15. Cohn. Die Refraction der Augen von 240 atropinisirten Dorfschulkindern, Arch. f. Ophth., xvii. p. 305, 1871.
16. Maklakoff. Examinations as to the occurrence of Myopia among School Children (Sitzbrt. d. phys. med. Ges. Moscow, 1871), Nagel's Jahresb., ii. p. 419.
17. Cohn. Die Augen der Schüler des Königl. Friedrichs-Gymnasiums und ihre Veränderungen im Laufe von $1\frac{1}{2}$ Jahren. Jahresbrt, d. K. Fr. Gymn. Breslau, 1872.

¹ The great majority of the eyes having exceptionally acute vision are hypermetropic, but, as shown, the majority of all eyes are H.


18. Krüger. Untersuchung der Augen der Schüler des Frankfurter Gymnasiums, Jahresbrt. ü. d. Verwaltung d. Medicinalwesens, Frankfurt-a-M., xv. p. 84, 1872.
19. Hugo v. Hoffmann. Augenuntersuchung in vier Weisbadener Schulen, Kl. Monatsbl. f. Augenh. xi. p. 269, 1873.
20. Aug. v. Reuss. Die Augen der Schüler des Leopoldstädter Communal-Real und Ober-Gymnasiums in Wien. Arch. f. Ophth., xxii. 1, p. 211, 1875.
21. A. Ott & E. Ritzmann. Bericht über die Untersuchung der Augen der Gymnasialisten in Schaffhausen, Correspondenz-Blatt f. schweizer Aerzte iv. 12, p. 321, 1874.
22. A. Ott. Der Refraktionszustand der Augen der Realschüler von Schaffhausen, Corrbl. f. schweizer Aerzte iv. 20, p. 583, 1874.
23. Cohn. Über die Augen von Greisen (Tagbl. d. Naturforscher-Versam. in Breslau, p. 105, 1874), Nagel, V. p. 222.
24. J. Gayat. Notes sur l'Hygiène oculaire dans les Écoles et dans la Ville de Lyon, Paris, 1874. Lyon Médical, xvi. pp. 12 and 87.
25. Max Burgl. Beiträge zur Aetiologie der Kurzsichtigkeit, Inaug. Diss. Munich, 1874.
26. H. Dor. Die Schule und die Kurzsichtigkeit, Rectoratsrede, Berne, 1874.
27. P. A. Callan. Examination of Colored School Children's Eyes," Am. Journ. Med. Sciences, lxxix. p. 331, 1875.
28. A. v. Reuss. Beiträge zur Kenntniss der Refraktions-veränderungen im jugendlichen Auge, Arch. f. Ophth., xxii. p. 211, 1875.
29. Pflüger. Untersuchung der Augen von 529 Lehrern, Kl. Monatsbl. f. Aughk., xliii. p. 324, 1875.
30. Max Conrad. Die Refraction von 3036 Augen von Schulkindern, 2d ed. Leipsic, 1876.
31. Hans Adler. Beobachtungen und Bemerkungen über das Sehen der Taubstummen, kl. Monatsbl., xiv. p. 65, 1876.
32. J. A. Spalding. Shortsight in Portland School Children, Trans. Maine Med. Assoc., 1874-76, p. 382, Portland, 1876.
33. Pflüger. Untersuchung der Augen der Lucerner Schuljugend, Arch. f. Ophth., xxii. 4, p. 63, 1876.
34. Emil Emmert. Ueber Refraktions- und Accommodations-Verhältnisse des Menschlichen Auges, Berne, 1876.
35. E. G. Loring. Is the Human Eye changing its Form under the Influence of Modern Education? New York, 1878. Trans. Internat. Med. Congress, Phila., 1876.
36. M. Scheiding. Untersuchungs-Resultate der Augen der Schüler an dem Gymnasium zu Erlangen, Inaug. Diss. Erlangen, 1876.
37. Oscar Koppe. Ophthalmoskopisch-ophthalmologische-Untersuchungen aus dem Dorpater Gymnasium und seiner Vorschule. Inaug. Diss. Dorpat, 1876.
38. L. Kotelmann. Die Augen der Gelehrtenschüler des Johanneums in Hamburg. Program der Schule. Hamburg, 1877.
39. L. Kotelmann. Cohn, Hygiene des Auges in den Schulen, Vienna and Leipsig, 1883.
40. A. Classen. Examination of the Eyes of the Pupils of the Realschule of the Johanneums in Hamburg, Nagel's Jahresbrt., viii. p. 371, 1877.
41. Wm. T. Bacon. Examination of the Eyes of School Children in Hartford, Conn. Private Communication.
42. Steven. Myopia in the Public Schools, Hartford Ev. Post, March 20, 1878.
43. Otto Becker. Das Auge und die Schule, VI. Univ. Lecture, Heidelberg, Centr. bl. f. pr. Augenh. p. 66, 1877.
44. C. R. Agnew. Nearsight in the Public Schools, N. Y. Med. Record, 34, 1877.
45. Hasket Derby. A Report on the Percentage of Nearsight in the Class of 1880 at Harvard College, Boston Med. and Surg. Jr., xviii. p. 337, 1877.
46. Niemann. Cohn, Die Verwechslung von Schschwäche und Kurzsichtigkeit, etc., Deutsch. med. Wochensch. 4, 1877.
47. H. Cohn. Die Augen der Uhrmacher, Goldarbeiter, Juweliere und Lithographen, Centblt. f. pr. Augenh., Apr. 1877.
48. H. Dor. Étude sur l'Hygiène Oculaire au Lycée de Lyon, Paris, 1878. Reprint, Lyon Médical.

49. A. Ott. Myopie und Schule, Corubl. f. schweiz. Aerzte, Nos. 15 and 16, 1878.
50. Seggel. Die Zunahme der Kurzsichtigkeit an den höheren Unterrichtsanstalten, Munich, 1878.
51. M. Reich. Einiges über die Augen der Armenier und Georgier in den Schulen von Tiflis, Arch. f. Ophth., xxiv. 3, p. 231, 1878.
52. Burchardt. Ueber die Verhütung der Kurzsichtigkeit, Deutsch. Med. Wochensch. No. 1, 1878, p. 6.
53. D. B. St. John Roosa. An examination under Atropine of the refractive state of eyes with normal vision, etc., Trans. Am. Ophth. Soc., ii. 4, p. 462, 1878.
54. —. Kurzsichtigkeit unter den Schülern in Hessen, Deutsche Z. f. Prakt. Medicin, No. 13, 1878, Nagel's Jahresbrt., ix. p. 418.
55. Haenel. Kurzsichtigkeit in Dresdener Schülern, Festschrift, Dresden, 1878.
56. Hasket Derby. Influence on the Refraction of Four Years of College Life, Trans. Am. Ophth. Soc., 1879.
57. Horstmann. Ueber Refractions-Verhältnisse von Kindern. Kl. Monatsbl. f. Augenh. Supl. 1879.
58. Eugen Netoliczka. Untersuchungen über Farbenblindheit und Kurzsichtigkeit, Graz, 1879. Abstract, xxviii. Jahresb. d. Steierm. Landes-Oberrealschule.
59. O. Just. Beiträge zur Statistik der Myopie und des Farbensinns, Arch. f. Augenh., viii. p. 191, 1879.
60. Nicati. La Myopie dans les Écoles de Marseille, Gaz. Hebdomadaire, p. 695, 1879.
61. Kotelmann. Die Augen von 9 Lappländern, etc., Berl. Klin. Wochensch., No. 47, 1879.
62. Kotelmann. Die Augen der Gymnasiasten und Realschüler in Wandsbeck, Program d. Gymn. 1879.
63. Priestley Smith. Shortsight in Relation to Education, Birmingham, 1880.
64. Ed. T. Ely. Ophthalmoscopic Observations upon the Refraction of the Eyes of newly-born Children, Archives of Ophthal., ix. p. 29, 1880.
65. Horstmann. Examination of the Eyes of newly-born Children, Centralblatt f. p. Augenh. Oct. 1889, p. 327.
66. Wm. S. Dennett. Report of Examination of Eyes of the Pupils in Schools of Hyde Park, Hyde Park, Mass., 1880.
67. B. Beheim Schwarzbach. Ueber Vorkommen und Behandlung von Augenkrank-
68.) heiten, in aussereuropäischen Ländern, Inaug. Diss. Würzburg, 1880.
69. E. Emmert. Auge und Schädel, Berlin, 1880.
70. B. Florschütz. Die Kurzsichtigkeit in den Coburger Schulen, Coburg, 1880.
71. Herzenstein. Die Untersuchung der Augen bei Schülern des Militär-gymnasiums zu Orel. Mil. Sanität Jaunr. No. 11. Centralbl. f. p. Augenh.
72. Schillbach. Examination of the Eyes of Pupils in Gymnasium of Jena, Jahresbrt. d. Gymn. Jena, 1879 and 1880.
73. E. Netoliczka. Untersuchungen über Farbenblindheit und Kurzsichtigkeit, ii. Graz, 1880.
74. L. Königstein. Untersuchungen an den Augen neugeborener Kinder, Med. Jahrbuch, i. p. 1, Vienna, 1881.
75. H. Cohn. Die Augen der Medicin-studirenden, Med. Jahrbuch. i. p. 21, Vienna, 1881.
76. A. v. Reuss. Augen-Untersuchungen an zwei Wiener Volksschulen, Wiener Med. Presse, Nos. 7 and 8, 1881.
77. M. Weber. Beitrag zur ophthalmologischen Schulhygiene, Inaug. Diss., Berlin, 1881.
78. S. D. Risley. Weak Eyes in Public Schools of Philadelphia, Trans. Penna. State Med. Soc., p. 789, 1881.
79. Badal. Examen des Yeux des deux cents Sourdes-Muettes de l'Institution Nationale de Bordeaux, Annales des Maladies de l'Oreille, du Larynx, etc., vii. 4, p. 198, 1881.
80. Q. C. Collard. De Oogen der Studenten aan te Rijks-Universiteit te Utrecht, Inaug. Diss., Utrecht, 1881.
81. Aug. Westphal. Untersuchung der Augen der Gymnasiasten, Jahresbrt. d. Gymnasium zu Schleiz, p. 26, Schleiz, 1881.
82. Eug. Netoliczka. Untersuchungen über Kurzsichtigkeit an den Grazer Mittelschulen, xxx. Jahresbrt. d. Steierm. Landes Oberrealschule.

83. Reich. Die Augen der Zöglinge des Tiflis'schen Militär-gymnasiums, Kaukas. med. Gesel. Centralbl. f. p. Augenh. p. 536, 1881.
84. Fox, L. Webster. Examination of Indians at the Government School in Carlisle, Pa., Phila. Med. Times, xii. 372, p. 346, 1882.
85. O. Paulsen. Ueber die Entstehung des Staphyloma posticum chorioideæ, Arch. f. Ophth., xxviii. p. 225, 1882.
86. Lyder Borthen. Refractions- und Farbenblindheits-Untersuchungen von Schulkindern, etc., Kl. Monatsbl. xx. p. 406, 1882.
87. P. Schubert. Ueber den Einfluss der Schiefschrift auf die Augen der Kindern Aerztl. Intelligenzbl., No. 21, 1882. Private Communication.
88. P. F. Roberts. Examen de la Vision practicado en las Escuelas publices de la Ciudad de Buenos Aires, Buenos Ayres, 1882. Cohn, Hygiene des Auges in den Schulen.
89. F. Erismann. Hygiene der Schule, Handb. d. Hygiene, etc., von Pettenkofer und Ziemschen, p. 2, 1882. Centbl. f. pr. Augenh.
90. Mittendorf. Einfluss der Civilization auf das menschliche Auge, etc., Verhand. d. Deutschen Gesellig-Wissenschaftlichen Vereins von New York, v. p. 30, 1882.
91. E. Nordenson. Recherches ophthalmométriques sur l'astigmatisme de la Cornée chez des écoliers de 7 à 20 ans, Extract, Annales d'Oculistique, livraison Mars-Avril, 1883, Ghent, 1883.
92. Wm. Manz. Ueber die Augen der Freiburger Schuljugend, Freiburg, 1883.
93. Berlin & Rembold. Untersuchungen über den Einfluss des Schreibens auf Auge und Körperhaltung des Schulkindes, second edition, Stuttgart, 1883.
94. Gaertner. Quoted by Berlin and Rembold, p. 46.
95. Hasket Derby. Influence on the Refraction of Four Years of College Life, Trans. Am. Ophth. Soc., iii. p. 456, 1883.
96. Dobrowolsky. The Eyes of the Pupils of the Ural High-school, Wratch No. 6, Nagel's Jahresbzt., xiv. p. 189.
97. Dürr. Die Refraktion von 414 Schülern nach Anwendung von Homatropin, Arch. f. Ophth., xxix. 1, p. 103.
98. Dürr. Ueber die Kurzsichtigkeit in den höheren Schulen Hannovers, Verhandl. u. Mittheil. d. Vereins. f. öff. Gesundheit, Hanover.
99. Wm. Hansen. Welche Refraction ist in 10 bis 15 Lebensjahre die vorherrschende? Kl. Monatsbl., xxi. 5, p. 196, May, 1883.
100. Reich. Nearsight in the Transcaucasian Girls' Institute, etc., Med. Sbornik. d. Kauk. Med. Ges. No. 34, Centralbl. f. Augenh., p. 483, 1883.
101. Motais. Hygiene de la vue chez les Typographes, Paris, 1883.
102. Schtschepotjeff. The Eyes of the Normal pupils of Astrachan, Tagb. d. Kasan. ärz. Vereins 13, Nagel, xvi. p. 579.
103. Schadow. Die Augen der Schulkinder Borkums, Kl. Monatsbl., xxi. p. 150, Apr. 1883.
104. Hadlow. Shortsight amongst Boys of the Greenwich Hospital School, Brit. Med. Jour., May 5, 1883.
105. M. Tscherning. Studien über die Aetiologie der Myopie, Arch. f. Ophth., xxix. 1. p. 201.
106. v. Reuss. Untersuchungen der Augen von Eisenbahn-Bediensteten auf Farbensinn & Refraction, Arch. f. Ophth., xxix. 2, p. 229.
107. Schleich. Die Augen 150 neugeborener Kinder ophthalmoskopisch untersucht. Mittheil. aus d. Ophth. Klinik zu Tübingen, ii. 1, p. 44, 1884.
108. H. Schäfer. Die Augen der Zöglinge der Taubstummenanstalt in Gerlachsheim (Grossh. Baden), Centralbl. f. Augenh., Mai, 1884.
109. Seggel. Ueber normale Sehschärfe und die Beziehung der Sehschärfe zu Refraction, Arch. f. Ophth., xxx. 2, p. 69, 1884.
110. Gustav Ulrich. Refraction und Papilla optica der Augen der Neugeborenen. Inaug. Diss. Königsberg, I. P. 1884.
111. J. Bjerrum. Trans. Internat. Med. Congress, Copenhagen, 1884. Quoted by Horstmann (vide 115).
112. Otto Beselin. Ueber Refraction und Grundlinie der Augen, etc., Arch. f. Augenh. xix. 4, p.
113. L. Kotelmann. Die Augen von 22 Kalmücken, Zeitsch. f. Ethnologie, xvi. 2, p. 77, 1884.
114. Seggel. Trans. Inter. Med. Congress, Copenhagen, 1884. Annales d'Oculistique.

115. Horstmann. Die Refractions-Verhältnisse des menschlichen Auges bis zum sechsten Lebensjahre, Ber. ü. d. xvi., Versammlung d. Ophth. Gesel., Heidelberg, 1884.
116. L. Webster Fox. Examination of the Eyes of Indian Girls from the Carlisle School. Private Communication.
117. H. Van Anrooy. De Oogen der Studenten aan te Rijks-Universiteit te Leiden, Inaug. Diss., Leyden, 1884.
118. M. Del Carlo & A. Pardini. Inchiasta Ottometrica nelle Scuole dell' Italia. Bollettino d'Oculistica, vi. 8 Apr. 1884.
119. M. Scellino. Inchiasta Ottometrica nelle Scuole dell' Italia, Bollettino d'Oculistica, vi. 9 and 11.
120. G. Masini. Idem. Bollettino d'Oculistica, vi. 10 bis.
121. G. Moyne. Idem. " " vi. 12 and 13 bis.
122. P. Brignone. Idem. " " vii. 2 and 3, 1884.
123. A. Hoffmann. Ueber Beziehungen der Refraction zu den Muskelverhältnissen des Auges. Inaug. Diss. Strassburg, 1884.
124. Schleich. Die Augen im Obergymnasium zu Tübingen. Private Communication.
125. Randall. A Study of the Eyes of Medical Students. Trans. Penna. State Med. Soc. 1885.¹

¹ The studies of Giraud-Teulon (Paris, Gaz. Hbd., p. 514, 1870), of Lopatin (Oljinski), of L. Zwingmann and of Buschbeck (Plauen, Vierteljrs. f. gericht. Med. u. öf. Sanitätsw. Jan. 1881), have been thus far inaccessible.



1806 Chestnut St. Phila


Dear Doctor

In the accompanying paper the attempt has been made to collect all the investigations bearing upon the subject of the statistics of refraction; and nearly twice as many studies have been brought together as in any previous summary of which the writer has knowledge. Complete success could not be expected, although four years have been spent in collecting data; and of the records available, many are very incomplete. Believing that the subject is worthy of very much more extended and closer study the writer deems it allowable to appeal to all interested in this question and in the host of others growing out of it, to kindly furnish him with any further data which may be available, with amplifications of reports which have not been published in full and most especially with corrections of anything which he has not accurately represented.

If you can kindly extend to him such aid
it will be gratefully acknowledged and
duly employed in the paper or papers in which
he proposes to continue this study. Criticism
as well as data will be gladly received, es-
pecially if any evidence of bias appears
in the treatment of the questions or facts un-
der consideration; and haste will be made
to correct any blunders which are pointed out.
It may be remarked that the writer has al-
ready gathered nearly one hundred of the
original records into his library, and has
access to many more; and he will continue
the effort to make the collection complete.
He hopes, therefore, that with the kind help
of the Profession his attempt to put these
statistics upon record in English will be
made a success, and engages to make the
fullest and most conscientiously ac-
curate use of all the data which he can
obtain.

John S. Billings M.D.
U.S.A.

Very respectfully
R. H. Randall.



THE MEDICAL NEWS.

A National Weekly Medical Periodical, containing 28-32 Quarto Pages in each Number. Published every Saturday.

Uniting the scholarship of a weekly magazine with the energy of a daily newspaper, THE MEDICAL NEWS renders a service of especial value to the profession. As a magazine, it contains *Original Articles* and *Lectures* by the ablest writers and teachers of the day, judicious *Excerpts* and *Translations* from the medical, pharmaceutical and scientific periodicals of the world, and numerous *Editorials* from the pens of a large and able Editorial Staff. As a newspaper, it maintains a corps of reporters and correspondents, in all the medical centres of the globe, and receives by mail and telegraph prompt and accurate reports of *Society Meetings* and other matters of immediate interest. It is thus enabled to present in each issue, without loss of time, the advance of knowledge attained in the medical sciences during the previous seven days. Ample space is devoted to *News Items* and *Correspondence* and to the weekly official list of changes in the *Army* and *Navy Medical Departments*. In typographical arrangement everything has been done to economize the time and to promote the comfort of its readers. The subscription price of THE NEWS has been fixed at the very low sum of Five Dollars per annum.

The American Journal of the Medical Sciences.

A Quarterly, Published on the first days of January, April, July and October, each Number containing over 800 large Octavo Pages, fully illustrated.

In his contribution to "A Century of American Medicine," published in 1876, Dr. John S. Billings, U. S. A., Librarian of the National Medical Library, Washington, thus graphically outlines the character and services of THE AMERICAN JOURNAL—"The ninety-seven volumes of this Journal need no eulogy. They contain many original papers of the highest value; nearly all the real criticisms and reviews which we possess; and such carefully prepared summaries of the progress of medical science, and abstracts and notices of foreign works, that from this file alone, were all other productions of the press for the last fifty years destroyed, it would be possible to reproduce the great majority of the real contributions of the world to medical science during that period."

This opinion of a man pre-eminently qualified to judge is corroborated by the great circle of readers of the Journal, which includes the thinkers of the profession in all parts of the world. During coming years the features of the Journal which have given unalloyed satisfaction to two generations of medical men, will be maintained in their vigorous maturity.

Together, THE AMERICAN JOURNAL and THE MEDICAL NEWS contain an equivalent of over 4000 octavo pages, in which duplication of matter is prevented by special management. They will be furnished together at a commutation rate of Nine Dollars per annum, in advance.

The safest mode of remittance is by Bank Check or Postal Money Order, drawn to the order of the undersigned; where these are not accessible, remittances may be made at our risk by forwarding in *Registered Letters*. Address,

LEA BROTHERS & CO.,
706 & 708 Sansom Street, Philadelphia.