

*With the Compliments*  
*Bulkley (D. D.)*  
*L. D. Bulkley M.D.*

# NEUROTIC ECZEMA

---

BY L. DUNCAN BULKLEY, A.M., M.D.

PHYSICIAN TO THE NEW YORK SKIN AND CANCER HOSPITAL, DERMATOLOGIST  
TO THE RANDALL'S ISLAND HOSPITAL, CONSULTING PHYSICIAN  
TO THE NEW YORK HOSPITAL, ETC.

---

*Reprinted from the Journal of the  
American Medical Association  
April 16th, 1898*

LIBRARY  
SURGEON GENERAL'S OFFICE

SEP. 1 - 1898

613

NEW YORK

1898

BY THE SAME AUTHOR.

---

FOURTH EDITION (READY VERY SHORTLY) MANUAL OF DISEASES OF THE SKIN. With Analysis of Twenty Thousand Consecutive Cases, and a Formulary. Price, \$1.25.

SYPHILIS IN THE INNOCENT (Syphilis Insontium) Clinically and Historically Considered, with a Plan for the Legal Control of the Disease. Price, \$3.50.

ACNE: Its Etiology, Pathology and Treatment. A Practical Treatise Based on the Study of One Thousand Five Hundred Cases of Sebaceous Disease. Price, \$2.00.

ECZEMA AND ITS MANAGEMENT. A Practical Treatise Based on a Study of Three Thousand Cases of the Disease. Second edition. Price, \$3.00.

NEUMANN'S HANDBOOK OF SKIN DISEASES. Translated, with Notes. Price, \$4.00.

ARCHIVES OF DERMATOLOGY. A Quarterly Journal of Skin and Venereal Diseases. Vols. I-VIII.; Vols. I-IV., \$3.00 each. Vols. V-VIII, \$4.00 each.

THE SKIN IN HEALTH AND DISEASE. Health Primer, 140 pages. Price, 50 cents.

THE USE AND VALUE OF ARSENIC IN DISEASES OF THE SKIN. Price, 50 cents.

ACNE AND ALOPECIA. Physician's Leisure Library. Price, 50 cents.

# NEUROTIC ECZEMA

---

BY L. DUNCAN BULKLEY, A.M., M.D.

PHYSICIAN TO THE NEW YORK SKIN AND CANCER HOSPITAL, DERMATOLOGIST  
TO THE RANDALL'S ISLAND HOSPITAL, CONSULTING PHYSICIAN  
TO THE NEW YORK HOSPITAL, ETC.

---

*Reprinted from the Journal of the  
American Medical Association  
April 16th, 1898*

LIBRARY  
SURGEON GENERAL'S OFFICE

SEP. 1 - 1898

613.

NEW YORK

1898



# NEUROTIC ECZEMA.

By L. DUNCAN BULKLEY, A.M., M.D.

Physician to the New York Skin and Cancer Hospital, Dermatologist to the Randall's Island Hospital, Consulting Physician New York Hospital, etc., New York, N. Y.

THE recognition and management of eczema must ever be an interesting and profitable study to the physician, because practically it forms a very large share of the cases of skin disease coming under his care. Even in dermatologic practice, where the rarer diseases and also the minor ailments of the skin contribute to make a considerable proportion of the whole number, it constitutes nearly one-third of all cases. In a recent analysis of 10,000 miscellaneous skin cases in the writer's private practice, there were 3,201 patients with eczema, they forming 32.01 per cent. of the whole number. In ordinary private general practice, where many infantile cases occur, this disease undoubtedly forms over one-half of all the cutaneous diseases coming for treatment.

Neurotic eczema is particularly interesting to the general physician, because if recognized and rightly treated not only are the results commonly very satisfactory, but the patient also receives much improvement in general health by the measures calculated to benefit the eczema. Neurotic eczema should always be recognized as the signal flag of danger and should never be passed over with attempts to gain relief from local treatment alone, which can only be of minor and temporary benefit.

The influence of the nervous system in the production of skin lesions of various kinds has long been recognized and is now abundantly established, both by pathological and clinical proof. Its influence in producing and prolonging eczema was very fully elaborated by the present writer\* some time ago, and reference will here be made to some of the evidence then collected, as well as to more recent clinical experience.

Neurotic eczema is seen in both sexes and in all ages, from the cradle to the grave, although it is much more common in certain periods of life than in others. In infancy it is frequently observed in connection with cutting of the teeth, fresh eruptions occurring as each new tooth presses on the swollen and tender

Presented to the Section on Cutaneous Medicine and Surgery, at the Forty-eighth Annual Meeting of the American Medical Association, at Philadelphia, Pa., June 1-4, 1897.

\*"On the Relation of Eczema to Disturbances of the Nervous System."—*Medical News*, Philadelphia, Jan. 31 and Feb. 7, 1891.

gum. In childhood it is less common, but may appear with each occurrence of nerve-strain, whether from overwork in school or over excitement in recreation. The same is true in youth, where the changes of puberty come in as a factor, especially in females. But its most frequent time of occurrence is during that active period of existence between 20 and 55 years of age, when the strain and burden of life falls heavily on so many and when the strongest constitutions too often show many evidences of breakdown, both in the nervous system and in other directions.

Something can be learned of this from the following tables. The first relates to nearly 6,000 cases of eczema, occurring among 20,000 miscellaneous skin cases in my public and private practice. The second table is constructed to show the percentage of eczema patients at different periods of life, as compared with the number of individuals alive at the same age, as shown by the life tables of insurance companies.

TABLE 1.—Ages of eczema patients.

|                      | MALES.  |         |        | FEMALES. |         |        | Totals. |
|----------------------|---------|---------|--------|----------|---------|--------|---------|
|                      | Private | Public. | Total. | Private  | Public. | Total. |         |
| Under 1 year.....    | 107     | 110     | 217    | 61       | 82      | 143    | 360     |
| 1 to 2 years.....    | 37      | 52      | 89     | 38       | 37      | 75     | 164     |
| 2 to 3 years.....    | 33      | 48      | 81     | 25       | 37      | 62     | 143     |
| 3 to 4 years.....    | 22      | 35      | 57     | 16       | 39      | 55     | 112     |
| 4 to 5 years.....    | 14      | 33      | 47     | 22       | 17      | 39     | 86      |
| Infantile.....       | 213     | 278     | 491    | 162      | 212     | 374    | 865     |
| 5 to 10 years.....   | 48      | 67      | 115    | 58       | 88      | 146    | 261     |
| 10 to 15 years.....  | 41      | 43      | 84     | 63       | 66      | 129    | 213     |
| 15 to 20 years.....  | 51      | 59      | 110    | 110      | 92      | 202    | 312     |
| 20 to 25 years.....  | 99      | 111     | 210    | 148      | 134     | 282    | 492     |
| 25 to 30 years.....  | 136     | 112     | 248    | 131      | 99      | 230    | 478     |
| 30 to 35 years.....  | 187     | 103     | 290    | 113      | 72      | 185    | 475     |
| 35 to 40 years.....  | 196     | 93      | 289    | 109      | 104     | 213    | 502     |
| 40 to 45 years.....  | 187     | 68      | 255    | 105      | 99      | 204    | 459     |
| 45 to 50 years.....  | 158     | 73      | 231    | 88       | 110     | 198    | 429     |
| 50 to 55 years.....  | 169     | 85      | 254    | 88       | 109     | 197    | 451     |
| 55 to 60 years.....  | 107     | 52      | 159    | 49       | 62      | 111    | 270     |
| 60 to 65 years.....  | 99      | 41      | 140    | 51       | 50      | 101    | 241     |
| 65 to 70 years.....  | 62      | 30      | 92     | 35       | 23      | 58     | 150     |
| 70 to 75 years.....  | 52      | 31      | 83     | 23       | 15      | 38     | 121     |
| 75 to 80 years.....  | 25      | 8       | 33     | 10       | 6       | 16     | 49      |
| 80 to 85 years.....  | 7       | 2       | 9      | 5        | 2       | 7      | 16      |
| 85 to 90 years.....  | 3       | 0       | 3      | 2        | 0       | 2      | 5       |
| 90 to 95 years.....  | 2       | 0       | 2      | 0        | 0       | 0      | 2       |
| 95 to 100 years..... | 0       | 0       | 0      | 0        | 1       | 1      | 1       |
| Unknown age.....     | 6       | 1       | 7      | 3        | 10      | 13     | 20      |
| Totals.....          | 1,848   | 1,257   | 3,105  | 1,353    | 1,354   | 2,707  | 5,812   |

TABLE 2.—Comparison of percentage of eczema patients at different ages, with the percentage of individuals alive at those ages as shown by the census.

| Ages.                  | Per cent. living, at all ages. | Per cent. of eczema patients. | Relation between the two. |
|------------------------|--------------------------------|-------------------------------|---------------------------|
| Under 1 year.....      | 2.2                            | 6.1                           | +3.9                      |
| 1 to 2 years.....      | 2.0                            | 2.8                           | -0.8                      |
| 2 to 3 years.....      | 1.9                            | 2.4                           | -0.5                      |
| 3 to 4 years.....      | 1.8                            | 1.9                           | +0.1                      |
| 4 to 5 years.....      | 1.8                            | 1.4                           | -0.4                      |
| Infantile.....         | 9.7                            | 14.8                          | +5.1                      |
| 5 to 10 years.....     | 8.7                            | 4.4                           | -4.3                      |
| 10 to 15 years.....    | 8.4                            | 3.6                           | -4.8                      |
| 15 to 20 years.....    | 8.2                            | 5.3                           | -2.9                      |
| 20 to 25 years.....    | 7.9                            | 8.4                           | +0.5                      |
| 25 to 30 years.....    | 7.5                            | 8.2                           | +0.7                      |
| 30 to 35 years.....    | 7.2                            | 8.1                           | +0.9                      |
| 35 to 40 years.....    | 6.7                            | 8.6                           | +1.9                      |
| 40 to 45 years.....    | 6.3                            | 7.8                           | +1.5                      |
| 45 to 50 years.....    | 5.9                            | 7.3                           | +1.4                      |
| 50 to 55 years.....    | 5.4                            | 7.7                           | +2.3                      |
| 55 to 60 years.....    | 4.8                            | 4.6                           | -0.2                      |
| 60 to 65 years.....    | 4.1                            | 4.1                           | ...                       |
| 65 to 70 years.....    | 3.3                            | 2.5                           | -0.8                      |
| 70 to 75 years.....    | 2.4                            | 2.0                           | -0.4                      |
| 75 to 80 years.....    | 1.5                            | 0.8                           | -0.7                      |
| 80 to 85 years.....    | 0.7                            | 0.2                           | -0.5                      |
| 85 to 90 years.....    | 0.2                            | 0.08                          | -0.1                      |
| 90 years and over..... | 0.07                           | 0.05                          | -0.0                      |

Examining these tables it is interesting to note the sudden increase in the frequency of eczema during the period from 20 to 25 years of age, and the continuance of the large number through those of 50 and 55 years in spite of the diminishing numbers alive at the advanced age. By the second table it will be learned that the greatest actual percentage of cases of eczema, in comparison with the individuals living, after the first year of life, was in the period between 50 and 55, where it was represented by + 2.3, while between 60 and 65 years the percentage was exactly equal; in the period between 10 and 15 years the disease is relatively the least frequent, the difference being represented by - 4.8.

It is also interesting to note the large increase in the number of cases in females, in private practice, between 20 and 25, where they form nearly 11 per cent. of all the females, whereas the males at this same period form only a little over 5 per cent. of the class. During the two decades between 20 and 40 years of age, when the strain of life is the greatest, there were no less than 1,947 cases of eczema, or almost exactly one-third of the entire number; this is the more remarkable considering the very

large number seen in the first years of life, and the diminished proportion of persons living during the later period, as shown in Table 2.

Various forms or phases of nerve disturbance are seen in connection with neurotic eczema, and they may be considered under the following heads: 1, Neurasthenia, or nerve exhaustion; 2, Nervous and mental shock; 3, Reflex phenomena; *a.* of internal origin; *b.* peripheral; 4, Neuroses; *a.* structural; *b.* functional.

1. *Neurasthenia or nerve exhaustion.*—First in importance, both from its frequency and its seriousness is nerve strain, which is a fertile cause of neurotic eczema. In many cases the direct causal connection may not always be immediately traceable, but watching the same patient for a length of time it will be found that with each period of nerve exhaustion, from varied causes, the eczema crops out. I cannot do better than to quote a few illustrative cases which I reported in the article previously alluded to.

An unmarried lady, aged 53 years, was the executive officer of a charitable institution, and her duties were arduous and exacting. On Tuesday of each week she was particularly tried on account of the weekly meeting and inspection of a board of lady visitors, and that night was often sleepless. Her eczema of the face and hands was always aggravated on the succeeding morning; indeed, the eruption, which would often yield very satisfactorily to treatment during the week would burst out afresh on the Wednesday morning succeeding the visit. This had been the case for some months before I saw her and was verified by myself again and again.

A prominent clergyman, aged 52 years, had for many years an eczema of the head and face, which was always greatly aggravated on Monday after the Sabbath's mental work and strain. This I observed on a number of occasions, the eruption greatly improving each week, until Monday, when he would present himself with an aggravation of the eruption. The same occurred sometimes during the week, whenever extra work, as a public address, was called for; the same has been repeatedly observed in other patients since.

In another case, that of a lady, age 22, any nervous excitement, or household disturbance and hysteric crying, would be followed by a fresh outbreak of eczema on the hands.

Since these cases were reported I have seen a number almost

as striking, and have frequently observed business and legal worry followed shortly by eczema. Mentioning the matter to an exceedingly intelligent gentleman, aged 47 years, whom I have attended in attacks of eczema for ten or more years, he remarked on the truth of the observation, and said, "you may quote me as a striking illustration of the effect of nervous strain in producing eczema again and again."

While writing this article, a gentleman aged 60, who has had eczema occasionally for fifteen or more years, also corroborated this statement very strongly.

Prolonged and severe household strain will produce the eruption, repeatedly, in women; and continued loss of sleep, often only from sickness in others, will now and again be followed by eczema. My notes show numerous cases where the nervous exhaustion following the grip caused the eruption to appear.

2. *Nervous and mental shock.*—The effect of nervous shock and violent emotions, such as grief, anger, etc., in the production of eczema has long been recognized by writers on dermatology, and the well authenticated instances of this on record are so numerous that it cannot be questioned that this form of nervous disturbance often operates either to produce the disease or to excite an attack in one susceptible to the same. A single very striking illustration may be cited from Leloir,\* a most careful authority.

"A young woman who never had any eruption previously, except a pustular disease of the scalp, at 2 years of age, but who was very nervous and impressionable, and who was in perfect health, suffered a mental shock as follows: Approaching her home she saw her little child leaning out of the window, and on the point of falling; she cried out, and the child being startled let go her hold and fell, strangely enough, directly into the mother's arms, so that it received no harm. But the mother lost consciousness, and was completely broken down by the shock, and within three days developed a sharp attack of eczema, beginning on the upper extremities and invading much of the surface. Some time afterward she had a second and a third attack, each time following mental disturbance."

---

\*Leloir: Des dermatoses par choc moral. Ann. de derm. et de syph., 1887, p. 367.

The same writer records several similar cases and Tommasoli\* reports the occurrence of eczema after long grief and sudden shock.

Mental and moral shock occurring in many different manners have been observed to be followed quickly by attacks of eczema. Thus Radouan† reports that cases occurred in immediate consequence of the siege and commune of Paris, in 1871, and I saw more than one case ascribable to the financial panic in Wall Street, known as "Black Friday," several years ago.

3. *Reflex phenomena, (a) of internal origin, (b) peripheral.*—The influence of reflex irritation in the production of disease has long been an established fact, although opinions have varied in regard to details and individual cases.

Many writers have related cases illustrative of the production of eczema by this means, and in certain directions it is of not very infrequent occurrence. This reflex irritation may come from within, from some other organ, or part of the system, or it may result from external irritation of the skin; in either case the eruption excited may cease with the removal of the cause, or may persist indefinitely.

(a) *Reflex eczema of internal origin.*—The most common illustration of this is the eczema observed in teething infants, where each accession of a tooth will cause a fresh outbreak of eruption on the cheeks, forehead, chin, or elsewhere. Another rather frequent form of reflex eczema is seen in that accompanying eye troubles. Many have reported cases of the eruption, which were cured by wearing proper glasses. I saw a very striking instance in a lady aged about 50 years, in whom attempts to use the eyes have repeatedly precipitated attacks of eczema of the lids and chin, also on the forehead. Intestinal irritation has been observed to produce attacks of eczema, and also tape-worm and oxyuris. Menstruation is likewise a frequent exciting cause, as observed by Danlos‡ and others, and as I have repeatedly witnessed.

(b) *Reflex eczema of peripheral origin.*—This form of disturbance is more difficult to make out, but a number of observations are on record where severe cutaneous irritation, as burns, etc., were followed by eczema. Scratching is a fertile cause of eczem-

\*Tommasoli: *Monatsh. fur pract. Dermatologie*, 1886, p. 432.

†Radouan: *Etude theor. et pract. sur l'Eczema*. Thèse de Paris, 1875, p. 41.

‡Danlos: *Etude sur la Menstruation*, etc., Paris, 1874.

atous eruption in those predisposed thereto. Not only does this result from the direct scratching of an affected part, but the irritation of a certain part by scratching or otherwise can excite a reflex irritation in a distant part, which may be followed by eczema in that locality.

4. *Eczema dependent upon neuroses, (a) structural, (b) functional.*—Eczema has been observed to follow structural nerve changes, as after injury, amputations, etc., and also in connection with functional neuroses, as neuralgia. It is not necessary to develop this part of the subject greatly, but I will mention one case where the connection of eczema and neuralgia was very marked.

Mrs. B., a widow, aged 30 years, was thrown on her own resources for the support of her mother and herself. She took to literary pursuits, and became the editor of a magazine requiring a large amount of personal work, she using the pen almost constantly. After some months she began to suffer from neuralgia of the right arm, and at the same time a papular eczema developed on the back of the arm and radial side of the forearm, attended with the most intense itching. When she ceased for a time from her great use of the hand the neuralgic pain lessened and the eczema diminished, but a return to her severe mental application would cause a return of both. This continued for some time, and treatment of either condition seemed to have very little effect while the arduous labors were persisted in, but both yielded when work was relaxed or suspended.

We have now reviewed some of the principal relationships of eczema with nerve disturbance, but the subject is a very much larger one than would appear from what has been said. The literature of it has already reached very great proportions and has hardly been alluded to. But enough has been said to show clearly the importance of the subject. We will now consider some of its more practical aspects.

Neurotic eczema does not differ very greatly from other forms of the eruption, but a trained eye can generally suspect the true character of the case.

The eruption is apt to come first upon the hands and face, less commonly on the feet. But from its starting point it may extend over large surfaces, and after scratching, or irritation by treatment, may present quite the features of gouty eczema.

Neurotic eczema on the hands is very apt to exhibit vesicles,

but on the adult face the eruption is quite as likely to assume and maintain the erythematous form, without vesicles, and often without moisture, unless scratched. The groups of lesions of neurotic eczema have a tendency to be pretty sharply defined, in more or less herpetic patches which may present mainly solid papules, or, when torn, a raw surface. The areas affected early in the eruption are not apt to be very large, but the eruption is composed of a number of patches of aggregated lesions.

Neurotic eczema is intensely itchy, and the patient will often scratch where there are no apparent lesions, and so develop the eruption in one place and another. The scratching indulged in is commonly of the most severe kind, it seeming almost impossible to reach the seat of the trouble until the deepest portions of the epidermis are reached. The spasms of itching are sometimes fearful and utterly uncontrollable. I remember the case of an elderly lady whose two daughters were not able, when the spasms seized her, though they used much force, to prevent her tearing her face and neck till the blood literally ran down, which I witnessed at my first visit.

**Treatment of Neurotic Eczema.**—This will often tax the patience and skill of the physician to the utmost, and the broadest principles of medical knowledge and judgment will often need to be put in operation and maintained, if the patient is to have great and permanent benefit. The treatment includes both constitutional and local measures; the former are essential, the latter are helpful.

*Constitutional treatment.*—Naturally every case requires to be studied and treated on its own merits; no two cases could be treated alike from beginning to end.

There is some danger of error, in approaching a case of neurotic eczema, in supposing that because of its nerve element it will require at once, and mainly, drugs which are known to have an influence on the nervous system. At times the treatment may be that of ordinary gouty eczema, for neurotic symptoms are exceedingly common in gouty subjects, and often largely dependent upon an existing gouty state, and unless that element is reached and rectified the progress will be slow. The strictest attention should be paid to the digestive system, the action of the bowels and kidneys, and above all to the diet; and this should be done, not only at the first visit, but on each subsequent consultation, and proper remedies and measures continually given.

But in all the treatment there is to be a continual regarding of the nervous system, and use of remedies and measures calculated to strengthen that. Arsenic undoubtedly finds an important place in the treatment of neurotic eczema, but should never be relied on alone, and I seldom administer it in the form of drops, the Fowler's solution so universally given by the profession for every case. Iron, nux vomica or strychnin, quinin, the phosphates, ergot, oils, and many other remedies may come into play in the management of the disease. In occasional cases much benefit will be obtained from digitalis in strengthening and regulating the capillary system, and so improving nerve and cellular nutrition, while other cases will require the sedative action of aconite in proper and free doses.

A few words may be added in regard to the attempt to give relief to the itching and securing sleep in neurotic eczema, by the administration of internal remedies, hypnotics. While the eruption is in an aggravated condition, it often seems almost impossible to effect much by this means. Opium and its preparations only aggravate the itching, and if sleep is secured it is of an entirely unrefreshing character. Codeia seems to be the least injurious of them all. Sulphonal and trional in large doses will sometimes be effective, but are often followed by further nervous exhaustion, in the end aggravating the complaint. Phenacetin in full doses, repeated in an hour or so, will sometimes prove most effective, and antifebrin, with hot water and a trifle of whisky, will often secure very refreshing sleep; in milder cases urethan in one gram doses is effective. Tincture of gelsemium, given in repeated and increasing doses, even every half hour, has in some instances proved most serviceable in my hands, as also cannabis indica. The bromids have relatively little effect, although when combined with a very small dose of morphia and aconite have at times proved very valuable. A large warm drink at bedtime will often aid these remedies and may be valuable alone.

The diet must always be carefully directed. Not only should ordinary rules be given as to the avoidance of unnecessary and injurious articles of food and pleasure, which by deranging the digestive system, can greatly hinder the progress of the case, but the diet should be so arranged that it shall furnish the best possible nerve nutriment, and in a form easily assimilable. For this purpose I always order an increase in the digestible fatty matter

and phosphates. Some caution may be required in regard to the former, but with a little care the amount of the fat of meats and oils, and also fresh butter, can be added to the dietary. Articles commonly called greasy, or where the fat is combined with starchy matter, as in pastry and fried articles, are to be avoided, but well browned bacon is often a very valuable addition to the meal. The phosphates are found abundantly in the preparations of whole wheat, such as crushed wheat, wheatena, wheatlets, wheat germs, Pettijohn's breakfast food, etc., as also in bread made from the whole wheat flour, some of which should be taken, if possible, three times daily. Fresh fish (not fried) is also serviceable.

But the one article which contains the elements necessary to nourish the nervous system and proves of the most signal advantage is milk, if properly taken. This should not be used at all with meals, nor after or in connection with the least particle of food, otherwise it undergoes caseation and may embarrass the liver, which is too often at fault in these cases. My plan is to have the milk taken warm, pure and alone, one hour before each meal, and also at bedtime, if sufficient time has elapsed for the stomach to be perfectly empty, which is at least four hours after a hearty meal. When the digestion is sluggish, and the milk meets the late products of digestion, it is curdled, and then its digestion requires some considerable time and we do not get the benefits desired. But if it can be taken on an absolutely empty stomach, when the alkaline tide has already set in, it is then absorbed almost immediately and affords a refreshment to the whole system, including the nerve elements, which can hardly be obtained in any other way, and which must be observed to be thoroughly appreciated. It is understood that this precludes the possibility of adding liquor or an egg to the milk, and especially should there never be a cracker or anything else eaten with or near it. When taken in the manner described not only does it not diminish the appetite, but increases it; whereas if by any chance the milk has become caseated in the stomach, the necessary digestion may, and generally does impair the appetite as well as disturb the liver action. This plan I have followed for many years, in hundreds of cases, and I am very positive of the vast benefit resulting, when it is strictly carried out.

*Local treatment.*—The indications for local treatment differ so materially in different cases that it would be impossible within

the limits of this paper to afford even the briefest outline of all that might be required; a few practical hints, however, may be given.

First, there is danger of doing too much in the attempt to get relief from the itching, which can seldom be obtained by local measures alone. The plan of treatment should be a soothing and protective one, and the mildest applications should be made first. Zinc ointment with 1 or 2 per cent. of carbolic acid or creosote, or with 5 to 10 per cent. of ichthyol, or tincture of camphor, is always a safe and generally beneficial dressing if well and thoroughly applied. But to be of service it should be kept thickly applied, spread on lint in most places, and bound on firmly. In the acutely inflamed and especially in the erythematous forms of the eruption, there is nothing better than the well-known calamine and zinc lotion, freely sopped on many times in the day. Ichthyol in watery solution, 10 to 20 per cent., freely bathed on the part, often gives great relief, or if the skin is too dry it may be used in oil in the same strength.

On more chronic patches the permanganate of potassium, 2 per cent. solution in water, painted over the part will sometimes arrest the itching very well; it may either be used alone, or the calamine and zinc lotion may be sopped on after the surface has dried, or an ointment may be applied if the skin is at all hard.

In the erythematous eczema of the face a tannin ointment, 1.95 to 3.9 grams, to the 31.2 grams, with 2 per cent. of carbolic acid is effective; a mixture of camphor, 1.95 grams to 31.2 grams of zinc ointment also forms a good antipruritic. In still more chronic states the tar and zinc ointment, as recommended by the present writer many years ago, when thickly spread on lint and bound on, will often also serve admirably to control the itching.

The use of very hot water for a brief application, followed by an appropriate ointment, should never be forgotten. In old cases of eczema of the scrotum the effect of this treatment is sometimes very remarkable. Menthol will often prove a valuable addition to ointments, in the strength of 2 to 4 per cent., with about half as much carbolic acid, which latter serves both to heighten the antipruritic effect and also to overcome some of the chilly sensation caused by the menthol.

In bringing this brief and rather imperfect study of neurotic eczema to a close, I wish to again emphasize what was intimated

early in my remarks; namely, that I do not regard eczema as a local condition or disease of the skin, but that it has constitutional and general relations which should always be considered and studied. The neurotic relations of eczema are especially interesting and important because of the excellent results which follow a careful and correct appreciation, interpretation and treatment of them; whereas, a neglect of this aspect of these cases will lead not only to their chronicity and rebelliousness, but also to an injury to the patient and further nerve exhaustion, whose ultimate result may be most deplorable.

4 East 37th Street.







# Syphilis in the Innocent

(SYPHILIS INSONTIUM)

CLINICALLY AND HISTORICALLY CONSIDERED

WITH

## A Plan for the Legal Control of the Disease

BY

L. DUNCAN BULKLEY, A.M., M.D.

### NOTICES OF THE PRESS.

- INTERNAT. JOUR. OF SURGERY, Sept., 1894.—“This work is the most important contribution to the literature of syphilis which has appeared within recent years.”
- BRITISH MEDICAL JOURNAL, Oct. 20, 1894.—“A veritable mine of information.”
- N. Y. MED. RECORD, April 21, 1894.—“The author has here gathered more facts and figures bearing upon the question of syphilis as a non-veneral disease than have ever before been published or in all probability attempted.”
- THE MED. NEWS, Aug. 11, 1894.—“The subject is an important one viewed from every light, and has been handled by the author in a masterly and practical manner.”
- ANNALES DE DERMAT. ET DE SYPH., April, 1894.—“The book must necessarily be consulted by everyone who has interest in the subject of extra-genital syphilis.”
- UNIV. MEDIC. MAGAZINE, Aug., 1894.—“The work is by all odds the best and most extended consideration of the subject in any language.”
- DUBLIN JOURNAL OF MED. SCIENCE, June, 1894.—“We are thankful for the book, and value it as a useful contribution to the study of syphilis.”
- CANADA LANCET, June, 1894.—“It is a work of great merit.”
- QUAR. ATLAS OF DERMATOLOGY, April, 1894.—“This is a book that no physician can be without. It is the only work on the subject, and is so complete that it leaves nothing to be desired. Buy the book and it will be money well spent.”
- VIRGINIA MED. MONTHLY, July, 1894.—“Every doctor should read this book.”
- JOURNAL OF CUTANEOUS AND GEN. URIN. DIS., June, 1894.—“One cannot help being impressed with the immense amount of research which must have been required to produce so complete an analytical bibliography as that presented in this work.”
- NEW ENGLAND MED. MONTHLY, April, 1894.—“The book is unique and absorbing from cover to cover.”
- THE MED. AND SURG. REPORTER, Dec. 8, 1894.—“That it is one of the most exhaustive works which has ever been made on any one medical subject is not too much to say of it.”
- AMER. JOURNAL OF MED. SCI., Dec., 1894.—“For the first time the subject of syphilis innocently acquired has been thoroughly studied, and the results of this investigation are perhaps as astonishing to the professional man as to the lay reader. \* \* \* \* The work is more than an able and laborious collection of statistics. It teaches a lesson so plain that no technical knowledge is needed to understand it. It should be read by every intelligent man and woman in this country. In the hands of moral reformers, the strivers after social purity, it would be a weapon more formidable than any with which they have ever been armed.”