

Valuable Information

for

Old and New Customers of

Porter's Pain King

How to get the Best Results
from the

Greatest Family Medicine

Since 1871

THE GEO. H. RUNDLE CO.

Sole Proprietors

PIQUA, OHIO, U. S. A.

To Our Friends and Customers, Users of Pain King

THIS pamphlet to the many friends of Porter's Pain King is written with the idea of giving them a better knowledge of the many things for which the medicine can be used.

We have been manufacturing this one article forty-five years and have thousands of customers who have used it since the first day it was made. We find, however, that even among our oldest customers there are a great many who have not the slightest idea of the many different uses of the medicine.

Please study carefully our circular which comes around the bottle; also read the following directions, and we believe you will find there is something about the uses of Pain King which you did not know before.

It is absolutely unnecessary for you to keep in the house different medicines for different ailments, for Pain King will do the work of each of them. For instance: Pain King can not be beaten when made into a cough syrup or salve and for all diseases or troubles you may have either in the home or at the barn among your stock.

Porter's Pain King contains no poison or dope, or anything injurious to the health. It is made only out of the purest drugs and herbs.

Do not be fooled by the big bottle other firms offer you. If you want a big bottle, pour a bottle of Pain King into a bottle twice as big, fill up with water and you will then have a better medicine than any other on the market, with the exception of pure Pain King.

HOUSEHOLD USE

INTERNAL AND EXTERNAL USE

Coughs and Asthma—Half pint of sugar syrup, one five-cent stick of licorice. Boil together, then add three teaspoonfuls of Pain King. For asthma add one-fourth pint of glycerine to the cough syrup.

Sore Throat—Mix two teaspoonfuls Pain King and one cup of sweet cream together. Use this as a gargle, applying medicine externally on the throat. Bandage the throat with greased flannel.

Salve for Cold on Lungs—Two ounces each of lard, rosin, beeswax or sheep tallow. Three tablespoonfuls of Pain King, one tablespoonful of turpentine. Mix thoroughly and apply to throat and chest; cover with flannel cloth.

Bronchitis or Croup—Heat equal parts of goose oil and Pain King and apply to throat and chest; cover with flannel cloth.

Hacking Cough or Tickling in the Throat—One tablespoonful of butter mixed with sugar, 6 drops of Pain King; hold portion of same in mouth until it melts.

Phthisic and Asthma—Take one teaspoonful in hot water; repeat the dose every hour until relieved; also bathe the throat and chest freely.

Lagrippe—Four teaspoonfuls of sugar, one teaspoonful of Pain King. Add one cup of hot milk. Stir until sugar is dissolved. This is one dose for adults; repeat every hour for three hours. Take one or two Porter's None Such Laxative Pills on retiring.

Lumbago—Bathe the back well with warm water and rub with rough towel, then bathe freely with Pain King.

Pleurisy—Bathe well with Pain King and saturate flannel with hot milk and Pain King and lay over the affected part.

Rheumatism—One tablespoonful of Pain King, three tablespoonfuls of sugar, one pint of warm water put in bottle and keep well corked. Take a tablespoonful from four to six times a day and bathe the affected part with pure Pain King.

Sprains—Bathe the part well with Pain King and repeat every hour until relieved.

Snake Bite—Keep the wound wet with the Pain King and take one teaspoonful in one-half cup warm sweetened water. Repeat every thirty minutes until relieved.

Chapped or Sore Hands—Equal parts of Pain King and glycerine. Rub hands thoroughly each night.

Cold Feet—Rub feet thoroughly with Pain King before going to bed.

Poison Ivy, Pimples, Sores, All Skin Diseases—Bathe freely with Pain King.

Poison Oak—Take equal parts of Pain King and sweet milk and apply to affected part.

Stomach and Kidneys—One part Pain King, sixteen parts black molasses, mixed thoroughly. Dose: A teaspoonful six times a day.

Burns, Scalds, Chapped Hands—For burns, Pain King and linseed oil equal parts. Mix thoroughly, applying with soft cloth. For chapped hands, one teaspoonful of Pain King to one-half pint water with a small portion of rose water.

Burns and Scalds—One part Pain King to one part sweet oil, or bathe freely with Pain King and bind up with cloth to keep out the air. Soak cloth with Pain King.

Cuts or Wounds—Wash wounds with warm water, then bind up with cloth soaked in Pain King. Do this as soon as possible after injury.

Cramps—One teaspoonful to one-half glass of warm water or milk. If not relieved in thirty minutes repeat dose until three doses have been taken.

Earache—A few drops of Pain King on cotton put in the ear will give relief.

Itch or Insect Bites—Bathe the part with Pain King and take internally one-half glass of milk or water to which has been added one-half teaspoonful of Pain King. Repeat in one or two hours if necessary.

Bee Sting—Bathe well with Pain King. Bind up the part with cloth soaked in Pain King.

Painful Menstruation—One teaspoonful of Pain King in half a glass of warm water. Apply externally, covering abdomen with warm flannel.

Cough and Croup—Make a syrup of sugar and water. To four tablespoonfuls of syrup add one of Pain King. Dose for adults, 12 or 15 drops every half hour. Children, 6 to 10 drops according to age. Bathe the throat and chest with Pain King. Keep the cough syrup bottled and well corked.

Tonsillitis—Bathe the throat with Pain King, make sacks about six inches square, fill them with bran and salt and a few hops. Apply to the throat hot. Repeat application of Pain King and the hot sacks every 10 or 15 minutes, until relieved. Very bad cases of tonsillitis have been cured in one day by the above treatment.

Felons—Put Pain King in a teacup or large mouthed bottle. Insert finger to first joint. Leave in the Pain King for 30 minutes.

Croup—One teaspoonful of melted lard and three teaspoonfuls of Pain King spread on cloth and fasten around the throat and over the breast. The lard prevents blistering.

Summer Complaint or Bowel Trouble—Make an ointment of one tablespoonful of lard, melted, and three tablespoonfuls of Pain King. Apply on bowels and cover with warm flannel cloth.

Rheumatism—Bathe affected part in warm water for five or ten minutes, then apply Pain King; rubbing in well. Cover with warm cloth saturated in Pain King. Repeat until relieved.

Corns—Bathe feet well; remove hard part of corn. Anoint corn night and morning with Pain King until it is entirely gone.

Skin Cancer—Anoint several times a day with Pain King until removed.

Frosted Feet—Apply freely to parts affected.

Fistula—Wash sore with warm water and castile soap, then apply Pain King three times a day until healed.

POULTRY

Chilled or Frozen Combs—Bathe combs with Pain King. Give 2 to 4 drops of Pain King in one and one-half cups clabber or water.

Cholera or Roup—Soak grain in Pain King water over night. Two tablespoonfuls of Pain King to quart of water. Also sponge head thoroughly with Pain King.

Lice—Apply liberal quantity either pure or slightly diluted with water. It will kill nits or eggs, so they will not mature.

Gapes—Dilute Pain King with water, dip feather in medicine and put down chicken's throat.

Limber Neck—Put pure Pain King in throat, being careful to get it below windpipe. Use feather or medicine dropper for this purpose.

HORSES

Colic—One-fourth bottle of Pain King in one pint of warm milk or water, or in very bad case, one-half bottle Pain King in one pint of warm milk. If not better in thirty minutes repeat the dose.

Sweeny and Ringbone—Apply Pain King clear and rub well.

Sunstroke—Four tablespoonfuls Pain King, one-half pint of warm water, add two tablespoonfuls of ginger, repeat dose every twenty minutes. Rub limbs and get the horse into shade if possible. Generally after second dose horse will get up.

Scours in Young Colts—One tablespoonful of Pain King, one-half pint warm milk and one raw egg. Repeat dose every hour until relieved.

Sore Shoulders—Wash thoroughly with castile soap and warm water, bathe with one-fourth Pain King to three-fourths sweet oil. Apply with sponge.

Sores—Saturate the sore with Pain King by applying it with a sponge or bunch of sheep wool for ten minutes.

Azatoria—Two ounces of spirits of nitre; in from ten to twenty minutes give two ounces Porter's Pain King and repeat the dose of Pain King in twenty minutes. Also bathe the horse over kidneys freely with the medicine, and in cold weather apply blankets until perspiration starts freely. Then give from one to two quarts of raw linseed oil.

Colds and Coughs—Mix one-third bottle Pain King with one pint molasses and after horse is through eating or drinking give about two tablespoonfuls of the mixture.

Cough Remedy—One-fourth of a bottle of Pain King; mix thoroughly with one quart of raw linseed oil. Dose, one to two tablespoonfuls on the grain.

Distemper—Pour some of the Pain King in hot water and hold under the horse's nostrils. Cover the head with a blanket so he will breathe the steam. This has cured many cases of distemper. Apply Pain King to throat with sponge or sheep wool.

Scratches, Calks, Kicks or any Flesh Wound of Horses—Melt one cup of mutton tallow, one cup of lard, add one tablespoonful of Pain King and enough starch to make a paste. This makes an excellent salve.

Pink Eye—Bathe eye with warm water. To two parts of warm milk or water add one part Pain King; inject into the eye with small syringe, repeat every four hours first 24 hours.

CATTLE

Bloat in Calves or Cattle—One-fourth to one-half bottle in one pint of warm milk or water. For calves, two tablespoonfuls in one-half pint of milk or water every twenty minutes.

Soreness of Udder—Equal parts of Pain King, linseed oil and turpentine; bathe freely with sponge or sheep wool.

Scours in Calves—One tablespoonful in one-half pint of warm milk and one raw egg, shake well. Given as a drench. Repeat every hour until you get results.

HOGS

Cholera or Indigestion—Drench the animal well with one tablespoonful of Pain King in one-half pint of water; repeat every two hours until relieved. Or, four tablespoonfuls Pain King, two tablespoonfuls salt, two tablespoonfuls pulverized charcoal, one quart of milk, given as a drench.