


UNITED STATES OF AMERICA




FOUNDED 1836

WASHINGTON, D.C.

AN
A T T E M P T
TO PROVE THAT *Stuart's*
D I G E S T I O N,
IN MAN,
DEPENDS ON THE
UNITED CAUSES
OF
SOLUTION AND FERMENTATION.




BY JOSEPH GLOVER,
OF CHARLESTON, SOUTH-CAROLINA,
MEMBER OF THE PHILADELPHIA MEDICAL AND CHEMICAL SOCIETIES.



_____ Tentanda via est, quà me quoq; possim.
Tollere humo. _____ VIR.



PHILADELPHIA:
PRINTED BY WAY & GROFF,
No. 48, North Third-street.
1800.



AN
INAUGURAL
EXPERIMENTAL INQUIRY,

FOR THE DEGREE OF

DOCTOR OF MEDICINE;

SUBMITTED TO THE

EXAMINATION

OF THE

REV. JOHN EWING, S. T. P. PROVOST;

THE

TRUSTEES & MEDICAL FACULTY,

OF THE

UNIVERSITY OF PENNSYLVANIA,

On the thirty-first of May 1800.

399021

TO
DOCTOR MATTHEW IRVINE,

PHYSICIAN OF CHARLESTON, SOUTH-CAROLINA,

WHO,

IN THE TWO-FOLD CHARACTER

OF A

FRIEND AND PRECEPTOR,

HAS THUS FAR GUIDED ME,

BY HIS

PATRONAGE AND CARE,

THROUGH AN

IMPORTANT BRANCH OF SCIENCE;

THIS

PUBLICATION

IS RESPECTFULLY INSCRIBED,

AS A SMALL TESTIMONY

OF

ESTEEM AND GRATITUDE,

BY HIS MUCH OBLIGED

FRIEND AND PUPIL,

J. GLOVER.

TO

DOCTOR MATTHEW IRVINE,

PROFESSOR OF CHEMISTRY, PENNSYLVANIA,

TO

BENJAMIN SMITH BARTON, M.D.

ON THE TWO-FOLD CHARACTER

OF THE SCIENCE OF MATHEMATICS, NATURAL HISTORY AND

AGRICULTURE, IN THE UNIVERSITY OF PENNSYLVANIA.

FRIEND AND PRECEPTOR,

THIS

DISSEMINATION

BY HIS

PATRONAGE AND CARE,

AS AN ACKNOWLEDGMENT

OF THE IMPORTANT BRANCH OF SCIENCE;

ATTENTION AND ADVICE

PUBLICATION

WITH WHICH HE HAS HONORED

IS RESPECTFULLY INSCRIBED
BY HIS MUCH OBLIGED FRIEND,

AS A SMALL TESTIMONY
J. GLOVER.

OF

ESTATE AND GRAYSTONE,

BY HIS MUCH OBLIGED

FRIEND AND Pupil,

J. GLOVER.

TO

BENJAMIN SMITH BARTON, M. D.

PROFESSOR OF MATERIA MEDICA, NATURAL HISTORY AND
BOTANY, IN THE UNIVERSITY OF PENNSYLVANIA,

THIS

DISSERTATION

IS LIKEWISE INSCRIBED,

AS AN ACKNOWLEDGMENT

OF THE

ATTENTION AND ADVICE

WITH WHICH HE HAS HONOURED

HIS MUCH OBLIGED FRIEND,

J. GLOVER.

P R E F A C E


BENJAMIN SMITH BARTON, M. D.

The difficulty of experimenting appears to be the reason why physiology has not kept pace with other branches of medical science. In most of these we may travel the smooth and delightful road; but the most trivial circumstances influence the result of the experiment. It is opposed to the reputation of the author to state, and on the reputation of the author to state, that the position is likewise regarded.

In the following pages it is contemplated to illustrate the process of digestion, a subject by no means generally understood. In doing this, I have endeavored to suggest such ideas on the subject as I have collected from the opinions of others or from such facts and experiments as I have myself ascertained. These have taught me the difficulty of explaining the phenomena which have been advanced; not do I expect that the one, which I have given, is too fond of reducing every operation of the animal economy to a single principle. Many physiologists have explained the process of digestion on some favorite theory, and thus by fixing a limit to the hand of nature, have left unaccounted for some of the most important phenomena.

ATTENTION AND ADVICE
WITH WHICH HE HAS HONORED

P R E F A C E.




THE difficulty of experimenting, appears to be the reason, why physiology has not kept pace with other branches of medical science. In most of these we may travel on a smooth and delightful road; but the most trivial circumstances influence the result of experiments; thus assertion is opposed to assertion, and on the reputation of the author rests the position.

In the following pages it is contemplated to investigate the process of digestion; a subject by no means perfectly understood. In doing this, I have endeavoured to arrange such ideas on the subject, as I have collected from the opinions of others, or from such facts and experiments as I have myself ascertained. These have taught me the difficulty of explaining the phenomena of digestion on most of the theories which have been advanced; nor do I expect that the one, which I have adopted, is void of imperfections. Too fond of reducing every operation of the animal œconomy to a single principle, many physiologists have explained the process of digestion on some favourite theory, and thus, by setting a limit to the hand of nature, have left unaccounted for some of the most important phenomena.

The result of my investigations, such as it is, circumstances have induced me to cast as my portion into the scale of science. Should it suggest an idea worthy the attention of the philosopher, I shall feel highly gratified; but, on the contrary, should my experiments prove inconclusive, and error stamp the reputation of my essay, still I shall be pleased with the reflection, that the most feeble attempt to elucidate this important branch of physiology, can be by no means injurious to science.


The haste, so unavoidably annexed to the short time allotted for this publication, will no doubt induce the reader to overlook the many inaccuracies of language, which I am fully aware are contained in the following pages. They may perhaps bring to his recollection, that sentence of Dr. Beddoes so expressive of the liberality of the philosopher, when he says, “we should set a due value on our present knowledge, though it be imperfect; and restrain those rude hands, that are ever ready to pluck up the tender plants of science, because they do not bear ripe fruit at a season when they can be only putting forth their blossoms.” Impressed with the generosity of these ideas, I submit my observations and experiments to the candid investigation of the philosopher, whose patronage alone stamps the merit of every youthful performance, gives activity to the mind, and sanctions future investigations.

AN
INAUGURAL
EXPERIMENTAL INQUIRY, &c.



SECTION I.

Observations on Digestion.



MAN, like every other being in nature, requires a continual and regular supply of food, for the several purposes of supporting life, of promoting the growth of his system until he arrives at maturity, and of forming new parts when such become necessary. Indeed, on contemplating the complicated structure of his frame, it is evident, that a continual loss of the solid and fluid parts, of which he is composed, must be the inevitable effect of every action or function of life which he performs.

Nature, always wise, to obviate this general waste of his system and continual tendency to decay, made it necessary that he should have some inducements to repair it, and thus be reminded of the connection which exists between aliment and life. Accordingly we find he is indued with the stimulus of hunger and thirst, which, together with the pleasure

ture he receives from gratifying those appetites, induce him to take into his stomach a certain quantity of matter to allay those disagreeable sensations. This matter includes, not only the several kinds of animal and vegetable substances which we denominate food, but also comprehends the fluids taken in with which they are diluted.

These having arrived at the stomach which is the great receptacle of his aliment, a greater or less length of time is requisite, according to circumstances, for them to undergo those processes which are essential to their assimilation being completed. In fact, I conceive it an impossibility to determine with any certainty, the exact time necessary for the digestive organs to perform their respective functions, as that will depend in a great measure on the strength of those organs, on the quality, quantity and manner in which the food is prepared, its previous mastication, and various other causes of which we are not always aware. The most prevalent opinion on this subject, is, that from about the third to the sixth hour after food is taken, it is discharged, through the pylorus, of a pultaceous consistence. There are however some extraordinary deviations from this allotted time, which cannot but convince us of the great uncertainty of calculations of this nature. We have on record instances related of substances remaining in the stomach indigested for months; and on the
contrary,

contrary, that in two hours after food was taken into the stomach, that organ was found empty.* These I conceive to be rare occurrences, neither do I believe it by any means common, even in the space of three hours, for the stomach to discharge its contents; as in a majority of mankind, I presume, a much longer time is requisite for food to undergo those changes, which are usually effected on it during its stay in that organ.

To our several kinds of aliment, different condiments are added by various nations, many of which merely gratify the palate, while others assist in promoting digestion. Among us at present sea salt is most universally used for this purpose.† Professor Chaptal tells us, that the acidulous tartrite of pot-ash is greatly consumed in the north of Europe, where it is used as a table seasoner; and Professor Barton has informed me, that the Creek Indians make use of hiccory ashes, and that in some of the southern states, the ashes of a particular kind of marsh-grass were formerly preferred for the same purpose.

Besides this difference in nations with respect to condiment, there is one of still more consequence, which is, their striking peculiarities in the choice of food. We find, that in Lapland, Iceland, Greenland,

* Vid. Haller's Element. Physiol. Tom. VI. page 281.

† According to the experiments of the celebrated Pringle, a small quantity of sea salt hastens putrefaction, while a larger quantity retards that process. *Diseases of the army, Appendix, paper 3. Exper. 25.*

Greenland, Norway, and other cold countries, the inhabitants live chiefly on fish and flesh;* while, on the contrary, we are informed of certain sects in India, who live almost, if not solely, on a vegetable diet. Both the Laplander and the Indian enjoy their health in these extremes; their habits, together with their climates, being better adapted to their respective modes of living; as the southern latitude, in which the latter resides, appears to prevent his subsisting for any length of time, on fish or flesh; while its stimulus is absolutely necessary to support the general waste of the system, to which the former is exposed from cold. A majority of mankind live in the medium of these extremes; experience having taught them, that a due proportion of animal and vegetable food, is the better adapted for their nourishment, the one counteracting the ill effects arising from the other.

Of all animals man appears to be the most omnivorous. Destined to range through every the most distant part of our globe, he is capable of accommodating himself to the food of every country. Certain other animals are likewise capable not only of changing their accustomed diet, but
sometimes

* This is sometimes (though more rarely) the case in southern latitudes. At Orange river, in Africa, Fordyce tells us, that the inhabitants live upon limpets, dead and putrid seals and whales, not tasting a particle of vegetable food of any kind, excepting aromatics.

sometimes acquire so vitiated a taste, as to refuse their former food. This was particularly the case with the wood pigeon of Spallanzani. This acute physiologist tells us, that by dint of hunger he brought this bird to relish flesh so well, that it refused every other kind of sustenance, even grain, of which it is naturally so greedy. Various other facts of a similar nature, are found on record. Von Troil informs us, that the Icelanders, when there is a scarcity of fodder, feed their cattle with *steenbitr*, (a kind of fish) which, together with the heads and bones of cod, is beaten small, and mixed with one quarter of chopped hay. He further adds, that the cattle are fond of this food, and yield a good deal of milk after having used it.*

Professor Barton, in his course on natural history, has likewise related a fact no less interesting. He has told us, that deer have been frequently observed to feed on dead fish, which had been washed up on the banks of the Susquehannah and other rivers. These are instances sufficient for my purpose, but many others of equal importance might be collected.

From the several kinds of aliment taken into the stomach, man is plentifully supplied with fluids, and from the component parts of these fluids, is the growth of his system and the solids of his body produced. This growth of his system and production of solids is induced, although he may sub-

sist

* Vid. Letters on Iceland, p. 133.

fiſt on very different kinds of food, as by the peculiar operation of his digeſtive organs, he is capable of aſſimilating, by certain proceſſes, matters taken from either the animal or vegetable kingdoms into a fluid *fui generis*. Theſe proceſſes of aſſimilation are comprehended in the term digeſtion; by it we are to underſtand, thoſe proceſſes which take place in the digeſtive organs of man, and by which his food is converted into laudable chyle.

The *modus operandi* of nature in this converſion of our food into chyle, has attracted the attention of philoſophers in every age, and various theories have been advanced to explain the phenomena which occur. With this view we find, that, the theories of *the heat of the ſtomach*, of *mechanical action*, of *fermentation*, of *ſolution* and others, have all been advocated by men, whoſe fame has added reputation to their opinions. But, as no one of theſe can excluſively account for all the phenomena of digeſtion, and as in the choice of a theory, the preference ſhould always be given to ſuch a one, as will explain to us the moſt of them; I am induced to adopt another, and attempt to prove the dependency of this important function, on the united cauſes of ſolution and fermentation.

Of the Heat of the Stomach.

The theory, of the heat of the ſtomach, was at one time ſo fashionable, that Profeſſor Blumenback

back tells us, instead of the term digestion, that of coction, was formerly used by the greater part of physiologists.* This opinion, however, I believe at present has but few advocates, as I presume no person will now contend that heat is the sole cause of digestion. This would be equally as incorrect to say, that it does not assist in promoting that process. While we refuse to admit, that heat is the sole efficient cause of digestion, we cannot but acknowledge its effect in expediting that process, as it has been long since made evident by experiment. It therefore only remains that we should shew from the situation of the stomach, that it is advantageously seated to be supplied with heat from its neighbouring parts, as we may easily conceive from contemplating their relative situations with respect to it. We find that its right side is covered by the thin edge of the liver; its left touches the spleen; that behind it is the seat of the pancreas, and immediately above it is the diaphragm; that the peritoneum lies before it, which, by the action of the abdominal muscles, gives it a motion diametrically opposite to that which it receives from the diaphragm in respiration; and that the aorta, the largest artery in the body, lies just behind it. All of these circumstances must tend to give it additional heat. Hence we may with propriety acknowledge the accuracy of Dr. Barry, when he tells us, that “the heat of
“ the

* Institut. Physiol. Vol. II. p. 23.

“ the stomach in a healthy man, is greater than the
 “ common heat of sun in a summer’s day.”*

Of Mechanical Action.

To disprove that the mechanical action of the stomach has any effect in promoting digestion, I need say but little. Facts speak for themselves. That accurate experimenter, the Abbe Spallanzani, has decidedly proven the very wonderful muscular action exerted in the stomachs of some animals; but his experiments likewise tend to shew, that no such action takes place in the human stomach. Having frequently swallowed wooden tubes during his experiments, which were made so thin as to be incapable of bearing the slightest pressure; he never, in a single instance, discovered one of them to be broken. In addition to this, he mentions the fact of cherries and grapes being voided entire, as I have myself frequently observed. He likewise relates an experiment with ripe grapes, which we all know to be incapable of bearing the least mechanical action, which appears to be directly in point. “ Of twenty-five,” says he, “ which I
 “ swallowed, eighteen were voided entire, of the
 “ other seven, the skins only appeared.”† These experiments alone, I deem sufficient to prove, that no triturating power is possessed by the human stomach,

* Vid. *Treat. on Digest.* p. 8.

† *Natural History*, Vol. I. p. 222.

mach, particularly as the fact of grapes being voided entire, must be notorious to every person who has attended to the subject. In short, I do believe, that the muscular fibres of the stomach, have no other effect on our food, than merely that of propelling it through the pylorus.

Of Fermentation.

While some physiologists of considerable reputation, have considered fermentation as quite sufficient to explain all the various phenomena of digestion, others of equal celebrity have contended, that no such process takes place in a healthy stomach. This diversity of opinion, I cannot attribute to motives of prepossession in favour of any particular theory, but would rather presume, it was the consequence of a supposition, that to admit the one to be a fact, would be a tacit acknowledgment that the other could not be true. This too I suppose is the reason, why even at the present time, those who have ascertained the solvent power of the gastric juice, will not admit that fermentation ever takes place in a healthy digestion. But this, perhaps though too common an error, is still one by no means the less prominent. Does not chemistry teach us, that nature frequently requires in her operations a multiplicity of causes to induce a single effect? Why then, if one of the causes, which have been advanced, is not sufficient to explain the phenomena which occur, should we attempt

attempt the establishment of another, equally inadequate to account for the wonderful effect of the conversion of our food into chyle?

The operations of nature are uniform, and frequently too deep for the shallow limits of human wisdom to demonstrate; but I think, when we shall have been more successful in our experiments, it will be found, that digestion depends on the combination of several causes, and that fermentation does certainly take place, as I shall endeavour to prove in a subsequent part of this essay.

Of Solution.

That solution does likewise take place in digestion, I do believe, as the experiments of many physiologists of considerable reputation tend to prove the position, and my own have induced me to embrace it. The opinion, however, is by no means novel. Barry informs us, that “Basil Valentine was the first among the chemists who supposed that animal digestion was owing to an acid dissolving menstruum.” That “Paracelsus received this opinion from him.” And that “Van Helmont carried it farther; and asserted, that the spleen prepared this menstruum, which was from thence conveyed into the stomach, by the *vasa brevia*.” Hence it appears, that the theory of solution has been long since favoured. Since when, very accurate experimenters have written, in hopes completely

pletely to decide the question in its favour. But, although they have most demonstratively proven the solvent power of the gastric juice, they have by no means shewn that fermentation does not likewise take place. For my own part, I do not hesitate in believing, that both solution and fermentation do take place in a healthy digestion; indeed I think, with correctness I may venture to assert, that in the human stomach, fermentation does as necessarily follow solution in the conversion of food into chyle, as thought succeeds impresson in the formation of ideas.

Food, in the first instance, is considerably attenuated, by the mastication which the rotatory motion of our jaws and pressure between our molares are so capable of giving it. Its particles being thus divided, are intimately mixed with the mucus of the mouth and saliva, after which it passes down into the stomach. This we may term a process preparatory to digestion. It is certainly one of much more consequence to the perfect digestion of our aliment, than is generally supposed; as it is evident those persons, who half chew or bolt their victuals, as it is called, are generally subject to all the numerous diseases arising from indigestion. Hence appears the necessity of persons being particular in the mastication of their food, as nothing scarcely can be of more injury to the constitution, than continually to supply the stomach with indigestible half-masticated food.

Something,

Something, in its effect very similar to mastication, is observable in domestic fowls. They, by a peculiar instinct, take into their gizzards, pebbles and gravel, which certainly serves in them every purpose, which teeth do in some other animals. I have made use of the word instinct through choice, because I cannot believe with Spallanzani, that they are picked up by mere accident, or through their ignorance in mistaking them for food. One fact appears to oppose his theory, which is, that those fowls which are kept on a gravelly soil are rarely if ever found to have a greater number of stones in their gizzards, than those raised where less gravel is present. Again, if they were picked up by accident, we should expect that they are not at all necessary to their health; whereas, the very reverse of this is the fact. A very respectable author, who investigated this subject with considerable success, by experimenting on chickens hatched with artificial heat, has given us the very best information I have perused. He says, “ I have hatched vast numbers, and frequently have given the chickens small feeds whole, taking care that they should have no stones. In this case the feed was hardly digested, and many of the chickens died. With the same treatment in every respect, others who had their feeds ground, or have been allowed to pick up stones, have none of them been lost.”*

This would appear to shew, that pebbles are essentially

* Ferdyce on Food, p. 24.

tially requisite to the healthy digestion of these animals. Indeed, the experience of many persons tends to prove this to be the case, as we often hear of their sending for gravel for their poultry, and when interrogated why they do this, they tell us, that without it their fowls grow poor, dwindle away, and sometimes die. Mr. John Hunter, commenting on this assertion of Spallanzani, that pebbles are picked up by birds through chance or ignorance, says with much humour that “it appears singular, that only those which have gizzards should be so stupid.”

The more freely food is masticated, and the more minutely it is divided; the less heavy does it lay on the stomach, and the more easy it is of digestion. The experiments of Spallanzani,* made on himself, prove the latter position; the former is obvious to common observation. Mastication not only promotes digestion, by minutely dividing the matters to be carried into the stomach; but likewise, by mixing them with saliva to form a pulpy mass, it involves a very considerable portion of atmospherical air, which, being rarefied by the heat of that organ, tends considerably to burst the several particles of food from each other. Here then we see that it indirectly promotes solution, for, as the several particles of food are separated from each other, in that proportion will solution be expedited; because, a greater number of surfaces will thereby be exposed

* Natural History. Vol. I. p. 224.


exposed to the action of the gastric juice, and of consequence the food will be the more speedily dissolved.

In the investigation of a subject like the present, it is necessary that we attend particularly to the causes which influence digestion, and judge of their effects, by comparing them to the effects of similar causes out of the body. And again, by accurately examining the products of digestion, and by comparing them to similar products out of the body, thereby investigate its causes. Thus, as all vegetable substances, capable of undergoing the vinous fermentation, contain the constituent principles of carbonic acid and of alcohol, and as these substances are obtained from such vegetables, the former during the fermentative process, and the latter after that process has ceased; it is evident, if such products are obtained from the digestion of food in the stomach, it must be the effect of a similar process. The same observations will apply to the other stages of fermentation.

In the following pages, I shall endeavour to relate in brief detail, such facts and experiments as have induced me to adopt the present theory. This will be done in three several sections. In the first I shall proceed to deliver a few observations on the matters which influence digestion. The second will contain an attempt to shew, how far solution is admissible in digestion. And in the third and last, I shall endeavour to relate, why it is presumable that fermentation does likewise take place.

SECTION.

SECTION II.



OF THE MATTERS WHICH INFLUENCE DIGESTION.

THE chief of these are, saliva, the gastric juice, the mucus of the primæ viæ, the bile, and the pancreatic juice. And,

1st. *Of the Saliva.*

By saliva we mean that colourless, glutinous, and resolvent fluid, which is secreted by the parotid, submaxillary, and sublingual glands. It is perfectly tasteless and inodorous in health, and consists of water holding a mucilage and certain salts in solution.

On being placed on the fire, its watery part speedily evaporates, leaving its salts in a state of crystallization, resembling two or three different kinds of salt. These are, according to Fordyce, sea-salt and sal-ammoniac, together with various other crystallizations of irregular forms; and agreeable to Plenck, sal-ammoniac and animal earth; the former he inferred from triturating quicklime with saliva; the latter from salival calculus and the products of fire.

Saliva from its sapid nature, gives an increased flavour to food. It tends to relieve thirst, by supplying the mouth and fauces with a sufficient quantity of moisture; and has a certain and evident effect in digestion. It does not however, possess a solvent

D

power

power, as has been asserted by some, at least it does not possess this property in a greater degree than simple water.

I put an ounce of pure saliva, and as much of simple water, into two separate phials; to each of these I added two drachms of roasted veal. These phials were placed uncorked in a sand bath, which was kept as near as possible to the heat of the human body. After suffering them to remain at rest for eight and forty hours, I decanted the water and saliva from each of the phials. The veal which had been immersed in water, and that which had been in saliva, were now placed separately on bibulous paper and as much of their moisture removed as possible. The one which had been in water, was now weighed; it had lost twenty-three grains of its weight. On weighing the other, it had lost no more. This would certainly seem to shew, that saliva has no solvent power, at least not out of the body.

Again, it has been asserted, that saliva corrodes copper and iron more speedily than simple water.* To ascertain this point, I made the following experiment:

Having placed twelve grains of sheet copper in a phial containing saliva, and the same quantity in another of equal size, containing water only; I put them both in a sand bath, which was generally about the temperature of one hundred of Fahrenheit.

* Vid. Haller's Element. Physiol. Tom. VI. p. 54. and Plenck on Hum. Fluids, p. 72.

heit. They were both kept in this situation for one week, at the expiration of which time, they were taken out, wiped dry, and weighed. I could not discover, that either one of them had lost more of its weight than the other. From this I presume we may conclude, that saliva has not a much greater effect, if any, in the corrosion of copper, than simple water.

Desirous of knowing what effect saliva has in digestion, and with a view of ascertaining whether it possessed a fermentative property or not, I exposed equal portions of leavened bread and roasted veal, in two separate flasks, to the heat of a sand bath, which I was careful to keep as near as possible to the temperature of the human body, or from ninety-six to ninety-eight of Fahrenheit. To one of these flasks, I had previously added eight ounces of water; to the other, seven ounces of water and one of saliva. The veal, to which saliva was added, I took the precaution of masticating, in order that it might be the more accurately blended with that fluid. In the space of six hours, the one containing saliva smelled a little sour, emitted air bubbles, and shewed evident marks of fermentation. Whereas, that process did not commence in the other, which contained water alone, until four hours after.

Professor Rush has long since made a similar and very decided experiment. "To elucidate the properties of saliva," says our author, "I placed
" mutton

“mutton and bread, of each two drachms, in two
 “glafs veffels. To one I added an ounce and a
 “half of faliva, to the other the fame quantity of
 “water, and placed both of them in a fand bath.
 “Five hours having elapsed, the mixture with fa-
 “liva began to ferment. In seven hours, it dif-
 “covered evident marks of acidity; whilst, in
 “that, to which water was added, fcarcely any
 “motion was perceptible. After twelve hours had
 “elapsed, the mixture with faliva emitted a putrid
 “smell; whilst the mixture with water remained
 “mild and inodorous to the twentieth hour.”†
 An experiment fomewhat fimilar, has likewise been
 made by Pringle, but not with exactly the fame
 fuccefs. He expofed two drachms of fresh meat
 and the fame quantity of bread, together with wa-
 ter and faliva, to the heat of a furnace, kept at
 the temperature of 100 of Fahrenheit. The mix-
 ture remained about two days, he fays, with fcarce
 any vifible fermentation; but on the third day that
 procefs became manifef.‡ This investigation, how-
 ever, does not appear to have been made fufficient-
 ly fair, for he tells us, that his experiment was
 made in a closed phial. Now we know that the
 prefence of vital air, is a circumftance effential to
 fermentation, and hence its exclusion must have
 been the caufe of that procefs being retarded. In

every

† Inaug. Differ. de Coctio. Cib. in Vent. Exper. V.

‡ Vid. Difcafes of the Army, Appendix, paper 4, Exper. 30.

every comparative experiment like this, every circumstance should surely be made as similar as possible, to those which occur in the living stomach, and of consequence there should be an admission of vital air, as this fluid is always involved in saliva, which is several times in the course of a single minute conveyed into the stomach.

We are informed, that some uncivilized nations, are so well aware of the fermentative property of saliva, that they prepare an intoxicating drink, by mixing saliva with certain vegetable substances. Plenck relates, that they prepare it from the chewed roots of the *jatropha manihot* or *cassada* and *piper methisticum*.

Whether saliva possesses a septic or antiseptic quality, has likewise been a subject of some controversy. To satisfy myself on this head, I placed equal quantities of roasted veal, in two separate phials of the same size. To one, I added an ounce of pure saliva, to the other, as much of simple water; these I placed, uncorked, in a sand bath, which I endeavoured to keep as near as possible to ninety-six of Fahrenheit. The heat of the bath, however, was sometimes as high as one hundred and ten, but never below fifty. In twenty-two hours, the one containing saliva emitted a putrid smell; the other only smelled sour. In eighteen hours after, I examined the phials again; they both smelled putrid; the one with saliva was the most
so,

fo, and likewise had changed its colour a little, which was not the case with the other.

From these experiments, I presume it appears, that saliva possesses the property of promoting both fermentation and putrefaction; and not that it promotes fermentation, and at the same time retards putrefaction, as has been supposed.

The quantity of twelve ounces of saliva, is generally supposed to be secreted by an healthy adult, in the space of twenty-four hours. At least this is according to the experiment of Nuck, whose assertion I believe continues to be considered as the most correct. The quantity however, is capable of being wonderfully increased by certain medicines and other stimulants. Indeed Haller speaks of fifteen ounces being effused in the space of thirty minutes.*

2d. *Of the Gastric Juice.*

The gastric juice in health, is a colourless and turbid fluid, void of both taste and smell; and I must add, bearing considerable resemblance to saliva. I do not believe, as some authors have supposed, that it is secreted from the small glands of the stomach, as it is inconceivable to me, how such minute glands could be the source of such large quantities of this fluid, as are at times secreted. I therefore shall prefer embracing the
theory

* Vid. Element. Physiolog. Tom. VI. p. 59.

theory of those, who suppose the immediate source of the gastric juice to be the extremities of the arteries of the stomach; for I can as easily conceive, in the wisdom of nature, that arteries may be expanded on the coats of the stomach in such manner as to perform secretions, as that they should be united together by ~~the~~ cellular membrane, in the form of glands for the same purpose.

Many persons are at present of opinion, that the gastric juice contains an acid. From the experiments of the Abbe Spallanzani, from those of his colleague Professor Scopoli, and indeed from those of many others, I think we have good reason to doubt of the presence of a sensible acid in the gastric juice.

Having obtained some pure gastric juice from my own stomach, I found it capable of coagulating milk very readily. This however I conceive as no proof of acidity being present, as I have coagulated it with a solution of fresh runnet, in which I could not detect the smallest particle of acid. Nothing can be more erroneous than the opinion which prevails among some persons, that acids alone have the power of curdling milk. The truth is the very reverse of this; for it is now known, that the heart, lungs, and even liver of a turkey have been discovered to possess this property.* It will surely not be said that they are likewise acid. Again, certain vegetables, as the galium luteum, or

ladies

* Vid. Spallanzani. Nat. Hist. Vol. I. p. 271.

ladies bed-straw, vaillantia cruciata or cross wort, rubia tinctorum or madder, carduus or thistle, cynara scolymus or artichoke, as well as many others, have been observed to have this effect. Indeed living fish have been observed to have this property; and Jacquin tells us,* that even lime-water produces an imperfect coagulation of milk.

Another fact, of no trivial import in the decision of this question, is related by Mr. John Hunter. This gentleman tells us, that “in the flink calf, “near the full time, there is no acid found in “the stomach; although the contents have the “same coagulating powers with those of animals “who have suck’d.”† Now, as this coagulating property is evidently communicated to the stomach by the gastric juice, and as an acid could not be detected in the stomachs of these young animals, although they possessed this coagulating property, I think it appears sufficiently clear, that an acid does not exist naturally in the gastric juice. This fact I consider as conclusive, at least in as far as it prevents our being deceived by acids which are evolved in digestion.

Haller likewise appears to have been fully persuaded, that pure gastric juice does not contain an acid, and has quoted the authority of at least a dozen persons to prove his assertion.‡ To determine
this

* Vid. Element. Chem. treat. de lacte.

† Vid. observations on the animal. œconom. p. 163.

‡ Vid. Element. physiol. Tom. VI. p. 143.

this point however more satisfactorily, as it is of such importance in the adoption of a theory of digestion, I made the following experiment.

Deeming it necessary to obtain the gastric juice perfectly free from any extraneous matter, and likewise from any acidity of a former digestion, I kept a cat fasting eight and forty hours, after which it was killed. I found no food in its stomach, and but a small quantity of gastric juice. This I submitted to the usual tests for detecting the presence of an acid, but could not discover any. Hence I have been induced to conclude, that the gastric juice does not contain a sensible acid, and that whenever an acid is present in that fluid, it must either be the effect of disease or proceed from the remains of some former food.

Barry says, “ that the humours which are contained in the stomachs of the most rapacious birds, fishes, and beasts of prey, have never an acid, but a saline taste.”* And from the chemical analysis of the gastric juice of the crow, by Professor Scopoli, we find he could not detect the presence of an acid in that fluid, but discovered it to be composed of, first, pure water; secondly, a saponaceous and gelatinous animal substance; thirdly, sal-ammoniac; and fourthly, an earthy matter similar, he says, to that which exists in all animal fluids.

We

* Treat. on Digestion. p. 22.

We come next to consider the solvent power of the gastric juice, which is the great basis on which the whole of the present favourite theory of digestion depends.

Spallanzani tells us, that this fluid in a dog, not only acts on bone, but even corroded the dense enamel of two dentes incisores taken from the upper jaw of a sheep.* And from experimenting on himself, he found he could digest not only “mucular fibres and membrane, but tendon, cartilage, and even bone itself, when not too hard.” From the excellent inaugural thesis of Dr. Stevens, we likewise learn, that various animal and vegetable substances, were dissolved by the gastric juice of the human stomach when inclosed in spheres. In like manner bone, and even ivory spheres were dissolved, which he had introduced into the stomach of a dog.

Since the publications of these two ingenious gentlemen, and particularly since the paper of Mr. John Hunter has appeared, further investigation of the solvent power of the gastric juice seemed unnecessary. This gentleman discovered, that in those persons who had died of violent deaths, the stomach itself was corroded and dissolved at its great extremity, and accounts for the stomach not being acted upon during life, on the theory of a *living principle*. Several facts tend to shew, that there is something innate in living matter, which resists the action

* Vid. Natural History, page 211.

tion of the gastric fluid. It is well known, for instance, that worms exist in the human stomach unaffected by digestion while living, but are speedily acted upon by the gastric juice as soon as they are deprived of life. To account for this fact, we must consider the gastric juice as a mere chemical solvent; then, by reflecting that all chemical solvents act by attraction, we may say, that the action of the vessels in living matter, is too great to be overcome by the attractive force of the gastric juice, and consequently their combination cannot take place. Another fact which shews, that the solvent power of the gastric juice is decidedly inert, as it respects some kinds of living matter, is, that there are several species of serpents, and particularly the rattle snake, who receive their young on the least alarm, down their throats into their stomachs, where they have been known to remain for three and four hours. Fifteen or twenty of these young rattle snakes, have been found in the stomach of an old one, and not in the least injured by the gastric juice.* “A polype” too “inserted into the stomach of another polype, continues to live as before.”†

Although convinced from these facts, of the inertness of the gastric juice on these animals, still I was not satisfied of its action on other living animals. I therefore determined on an experiment with the gastric juice of a dog. For this purpose I kept a
dog

* Professor Barton's Lectures on Natural History.

† Spallanzani's Natural History, Vol. I. p. 111.

dog fasting twenty-four hours, at the expiration of which time I obliged him to swallow a number of small sponges. As his stomach retained these very readily and without any apparent inconvenience, I suffered him to remain at rest for three hours, immediately after which he was killed. From these sponges I obtained about half an ounce of very pure gastric juice. In it I immersed a common earth-worm. The animal writhed about, and shewed symptoms of great distress. I corked the phial, and being at some distance from home, walked with it in my pocket to my residence. At this time I examined it again; exactly half an hour had elapsed from the time of my placing it in the phial. The animal shewed no symptom of life, and on taking it out of the phial, I found on its body evident marks of violent inflammation. Suspecting this, however, to be the consequence of the heat of my pocket, which (as it was in the month of April) I believe to have been about 80 of Fahrenheit, I repeated the experiment. The animal after being immersed in the gastric juice, at the temperature of 70 of Fahrenheit, for fifty minutes, shewed no signs of life; but there was no inflammation evident, as in the former. Knowing, however, that these animals live in a temperature still lower than this, I determined again to repeat the experiment. To effect this I placed an earth-worm in the phial of gastric juice as before, and covered it over with the
same

same food and in the same spot from which the animal was taken. Fahrenheit's thermometer stood in the turf at 60. In an hour I examined the phial, took out the worm, and found it lively and not in the least injured; but on increasing the temperature to 70, it died as the last. Hence we may infer, that this animal also resists the action of the gastric juice while living and at its own temperature. How far this will be found to be the case with other animals, I am at present unprepared to decide. I am aware that, according to the experiments of Dr. Stevens, a living leech, which some persons have supposed to be an animal destitute of pores and capable of sustaining a degree of heat equal to the human temperature, is digested by the human stomach.* But, that they are capable of sustaining this degree of heat, I am not satisfied; at least, of this we may be certain, that the temperature, at which they usually live, is not by any means as high. I had rather presume, that as in my experiments, the animal in this instance, had died previous to its being acted on by the gastric juice. Plot does indeed tell us, that he has seen the eyes of a carp and the nose of a roach, which were taken out of a jack-fish, digested, while they were yet alive.† And Cornelius is likewise said to have found a snake half digested in a bird's stomach, while life in that animal was still perceptible.

In

* Vid. his Inaug. Differ. pub. at Edin. in 1777.

† Vid. Plot's Staffordshire. Ch. 7. Sect. 37.

In all experiments with a view to ascertain the action of the gastric juice on different kinds of living animals, to proceed fairly I think we should keep this fluid exactly at the natural temperature of the animal immersed in it, otherwise it may be destroyed, either by an excess, or too great abstraction of heat. The gastric juice of different animals too, requires as different temperatures to promote their action. Thus we learn that this fluid in amphibious animals and in fishes, acts on aliment at the common temperature of the atmosphere; but in most animals of the class mammalia, a higher temperature is requisite. The gastric juice of different animals varies in another respect still more considerably. Thus, in some animals, we see that it appears destined to act on animal food only; in others, on vegetables alone; and in others again, on both.

The gastric juice of man acts on antimony and copper; the former I relate on the authority of Professor Chaptal,† the latter on that of Professor Barton. This last gentleman mentions, in his lectures, the case of a child, who was much indisposed and salivated, in consequence of swallowing a cent; the cent when voided, was examined and found to be sensibly diminished.

According to Jacquin, Spallanzani and others, the gastric juice of itself has little tendency to either

† Vid. *Elemen. Chem.* Vol. II. p. 260.

fermentation or putrefaction ; but when mixed with other substances, its effect is rather to retard those processes.

From the numberless small arteries of the stomach, we may presume the quantity of gastric juice secreted to be very great ; but like other secretions it is increased in quantity by stimuli, and particularly by the stimulus of food.

3d. *Of the Mucus of the Primæ Viæ.*

This mucus is found on the internal surface of the stomach and intestines, and covers it very completely. It has the appearance, consistence, and properties of mucus in other parts of the body. It is secreted from small glands, which are situated under the villous coat of the primæ viæ. The quantity secreted appears to be very great.

Its use seems to be that of lubricating the surface of the primæ viæ, in order to facilitate the passage of their contents. It must likewise defend the internal surface of the stomach and intestines, from the action of the gastric juice, and from the acridity of bile when regurgitated.

To ascertain whether this mucus has any effect, either in promoting or protracting the process of fermentation out of the body, I placed equal portions of leavened bread in two flasks, each containing eight ounces of water. Reserving one of these as a criterion, I added to the other about four drachms

drachms of this mucus, which I had procured from the stomach of a subject, who had died of a fall. Both flasks were placed in a sand bath, which was kept as nearly as possible at the temperature of the human body. In seven hours, the one containing mucus emitted air bubbles and smelled sour, whereas the one without mucus had no appearance of fermentation for several hours after.

4th. *Of the Bile.*

The name of bile, has been uniformly given to a peculiar fluid, exclusively found in animal bodies, and which is secreted from a gland of a singular structure, called the liver. It is more or less of a yellowish-green colour, of a disagreeable bitter taste, of a thicker consistence and more plastic than saliva, of a singular aromatic smell when evaporated, which has been compared to that of musk, and which is by some thought agreeable.

Bile differs from all other secreted fluids in the body, in this, that it is not like them, separated from florid arterial blood, but is secreted from the dark coloured blood of the vena portarum, which is nothing more than a large vessel, made up by the concurrence of all the veins of the viscera of the abdomen, (except those of the liver,) which empty their blood into it for the purpose of secretion. Now a plentiful supply of blood to a gland, being essential to the secretion of a fluid, and this blood being

being conveyed to the liver by the vena portarum, it has, and I think with propriety, been said, to perform the office of an artery. It ramifies every where throughout the substance of the liver, and terminates in two very different kinds of vessels. The one returns the blood, which is no longer fit for secretion, again into the general circulation; the others are the secretory vessels. In the former case, the extremities of the vena portarum inosculate with those of the hepatic veins, and thus the blood is returned to the inferior or ascending cava. In the latter, the secreting vessels soon terminate in the pori biliarii, by the union of which, in their passage out of the liver, trunks of a larger size are gradually formed, till at length they terminate in one of considerable magnitude, known by the name of the hepatic duct. This duct in its turn, soon terminates in another, which has gotten the name of the ductus communis choledochus, in consequence of its being the common duct of the liver and gall-bladder, through which bile is continually distilling into the duodenum, in which intestine this duct ends. Just however where the hepatic duct ends, and where the ductus communis choledochus begins, another duct arises, which extends to the gall-bladder, from whence it has received the name of ductus cysticus.

The secretion of bile, is a subject as yet involved in considerable obscurity. While some physiolo-

gists have considered it as the effect of a peculiar secretion in the liver, others of equal reputation, have asserted, that it is found formally in the blood. The correctness of these assertions are only to be determined by experiment, as therefore I cannot do this at present, I shall not venture a conjecture on the subject.

The bile being once secreted, it is received by the small branches of the hepatic duct called *pori biliarii*, from whence it passes into larger branches, till at length it gradually arrives at the great trunk of the hepatic duct; from thence it passes through the *ductus communis choledochus*, and is finally discharged into the duodenum. This is the most common course of the bile, but if from any spasmodic affection or morbid distention of the duodenum, or from any obstruction in the *ductus communis choledochus* its passage should be prevented, a retrograde motion of this fluid is the consequence. In this case it regurgitates through the cystic duct and finds its way into the gall-bladder, which is a very convenient receptacle, destined to prevent the hepatic duct from being surcharged with bile, in cases either of obstruction or of too great secretion.

Cystic bile differs, it appears, from hepatic, in this, that its consistence is more grumous, it is of a darker hue, and has a more pungent bitter taste. They do not differ in their constituent parts, but only in the proportion of those parts. In fact

I do

I do believe, that the only material difference which exists between them, depends on the more aqueous part of the cystic bile being absorbed, from its confinement and stagnation in the gall-bladder. Two causes concur to promote the discharge of cystic bile. The one is, the pressure which the gall-bladder receives from the neighbouring parts, and particularly from the stomach when distended with food; the other, that either from the acrimony which bile acquires by stagnation, or from the mere stimulus of distention, a contraction of the muscular coat of the gall-bladder will be induced, and its contents will thus be readily propelled into the duodenum.

From the experiments of the most accurate chemists, the constituent parts of the bile appear to be, 1. A coagulable lymph, 2. A resinous matter, 3. Animal gluten, 4. Soda, and 5. A colouring matter, which is believed to be iron. Neither of these component parts of the bile is present in a perfectly free and disengaged state; they are all so combined as to form an apparently homogeneous fluid. Bile, without the assistance of an intermedium, is not miscible, as far as I know, with any volatile or fixed oil, with animal fat, nor yet with butter or any other oleaginous substance. Neither does it appear to render these substances miscible with water, although perfectly so itself. I have seen some of the bile of an ox, which had been mixed in a phial with an equal quantity of olive oil, and

which

which after having been kept for more than twelve months and frequently agitated, had not the least disposition to unite with it. The idea of its rendering oils miscible with water, appears to have arisen from its being long since used in the cleaning of stuffs, and hence it has been supposed to act chemically and in the same manner in which soap does. It does I believe act chemically, but still not in exactly the same manner in which soap does. Soap is said to have an attraction for both oil and water, and thus renders them miscible; whereas, bile by having a greater attraction for the stuff to be cleaned, than oil has, only tends to displace it, and it is in this way that I believe it always acts, when used to remove spots of oil, or other greasy matter, from substances to which they are adherent.

The use of the bile in the animal œconomy is evidently material, and I may add, it is singularly important in the process of digestion. I do not however believe, that it has any effect in this process while going on in the stomach, as its presence in this organ is the consequence of regurgitation, and is without doubt morbid. This I infer, first, from the sickness of stomach, vomiting, vertigo and other symptoms of great morbid action, which attend its presence in that organ. And secondly, from these symptoms being speedily relieved, by such medicines as effect its discharge.

Doctor Monro having caught several frogs, at different times killed three of them, and as speedily

as possible emptied the contents of their gall-bladders into the stomach of another. The consequence of this was, that the animal died shortly after.* I do not attempt to prove any thing more from this, than that there is somewhat deleterious in the bile of these animals, particularly as we are aware that the bile of other animals have been taken not only with impunity, but even with advantage. I have poured the recent cystic bile of a cat down the throat of a puppy, without his suffering the least inconvenience from it; and I have seen a black servant whose taste had become so vitiated, after having accustomed himself to take the bile of an ox frequently as a stomachic, that he became fond of it, and so far from its proving injurious to him, he always thought himself much better after using it.

Bile neutralizes both the vegetable and mineral acids and is itself decomposed by them. In the duodenum, by mixing with the chymous mass discharged from the stomach, it first begins to separate the chyle from it, and being itself decomposed, its more aqueous part unites with the chyle, while its resinous adheres to the fæces giving them their natural yellow colour; thus the former is prepared to undergo the round of the circulation, the latter to be discharged from the body.

“ A

* Dr. Wistar's Lectures.

“ A further use of the bile,” says an admired author, “ is, to evolve and exterminate from the alimentary canal, the fixed air, which had been hitherto confined among the chymous mafs.”* Again, it prevents the too great accumulation of mucus in the intestines, and by its stimulus increases their peristaltic motion. Hence it is that biliary obstructions are generally accompanied with costiveness, and sometimes with a discharge of mucus.

The property of being both powerfully antiseptic and antizeumic, is likewise said to be possessed in an especial manner by the bile. Maclurg tells us, that “ this fluid after having remained to his knowledge, for three days in a dead body, and although when the gall-bladder was taken out, there was a very offensive smell in all the abdominal viscera ; yet this fluid, being poured into a phial, and closely stopped, acquired a sweet smell, which continued some days before the putrid fetor began.”† This property of bile has been supposed to be intimately connected with its bitterness. Knowing, however, that bile is secreted from the dark coloured blood of the vena portarum, (which is entirely free from any changes effected by air, through the medium of the lungs, and in fact possessing all the characters of common venous blood,) does it not appear probable, that its antiseptic quality depends on its attraction for, and consequent absorption

* Blumenbach.

† Vid. Treat. on the Human Bile. p. 76.

absorption of, oxygene. The opinion that the blood, by its circulation through the abdominal viscera, receives a putrescent tendency, appears to be erroneous, as it is deprived of its oxygene and consequently becomes antiputrescent; for bodies can only become putrid by the absorption of this gas.

The difference which exists between the blood of a foetus, which is appropriated to the secretion of bile, and that of an adult, appears to be worthy of some attention. The difference is briefly this, the blood, from which the bile of a foetus, is secreted, partakes more of the quality of arterial blood than that of an adult; this likewise makes a proportionate variation in the properties of the bile, and consequently that of a foetus is of a more putrescent nature, or in other words, has less tendency to resist putrefaction.

The quantity of this fluid secreted seems evidently to be great, particularly when we consider the vast apparatus of its secretory organ, and the quantity of blood conveyed to it for this purpose. Indeed Dr. Monro tells us, that four ounces of cystic bile have flowed through an ulcer of the side daily.*

5th. *Of the Pancreatic Juice.*

The juice denominated pancreatic, is a limpid fluid, which bears a greater resemblance to saliva

* Vid. System of Anatomy. Vol. II. p. 389.

saliva than to any other fluid in the human body. It is secreted from a long and flat gland of the conglomerate kind, which lies under the stomach, and between the liver and the spleen, and which is known to anatomists by the name of the pancreas. The situation of this gland in the abdomen is transverse, being in the duplicature of the posterior portion of the mesocolon. It is found not only in man, but in most other animals, in quadrupeds, in birds, and in many fishes.

The external appearance of the pancreas, is that of one uniform substance, with its surface somewhat uneven from a considerable number of small convexities, and resembling very much in its structure the salivary glands. In the centre of the breadth of this gland, we find its great duct running in a longitudinal direction, and into which several smaller ones empty themselves on each side, like so many minute branches inosculating with one parent stalk. It has very properly gotten the name of the pancreatic duct, and opens generally in common with the ductus communis choledochus into the duodenum, for the purpose of emulging its contents. This however is not always found to be the case, as it sometimes opens by a separate duct into that intestine.

An exact analysis of the pancreatic juice, as far as I know, has never yet been made; but, like most of the fluids of the human body, it is found

to contain common falt and fal-ammoniac. The difficulty of procuring a fufficient quantity of this fluid, I believe, is the reafon, why its analyfis has not hitherto been much attended to. We are directed to obtain it by inserting a fmall tube, to which a phial is appended, into the pancreatic duct of a living animal; but this cannot be fo readily accomplifhed, as we may at firft fight imagine.

The ufe of this fluid is not perfectly underftood. It is generally believed to have the effect of diluting the chymous mafs, after its having paffed into the duodenum from the ftomach, and to affimilate it to an animal nature. Likewise to dilute and attenuate the bile, which is fometimes too vifcid and acrid.


From analogy the quantity of this fluid fecreted appears to be very great; as the pancreas is no lefs than three times as large as the falivary glands, and has every circumftance as favourable for the fecretion of its fluid. Like other glands in the body, its fecretion is increafed by ftimulants, which no doubt makes a very confiderable variation in the quantity fecreted at different times. The preffure which this gland receives from the ftomach when diftended with food; the irritation of the chyme in its paffage into the duodenum; and even that of the bile itfelf, tends to promote the difcharge of its juice. Like the bile, Haller fays,* it is capable of being regurgitated into the ftomach.

G

SECTION

* Element. Phyfiol. Tom. VI. page 309.

SECTION III.



HOW FAR SOLUTION IS ADMISSIBLE IN DIGESTION.

TO assert that solution does not take place in digestion, would be to deny every thing like testimony in medicine. My attempt will only be to shew, why it is not probable that it can be the sole efficient cause of that process, and how it should be considered as tending to promote it.

By solution we can comprehend nothing more, than so minute a division of the particles of any matter, as to render that matter capable of being dissolved in a fluid; and this too, without effecting any change of its component parts; for no possible division be it ever so minute can have this effect. Thus, the component parts of water are the same, whether it be in the state of ice or of vapour. If this be a correct definition of what we understand by the term solution, a solvent in the stomach can have no other effect on our food, than merely that of separating it into very minute parts or particles; but this is very far from being all which takes place in digestion. Such a change must be effected on our aliment, for the due nourishment and support of our systems, as to convert it into that mild and bland fluid which we denominate chyle. This fluid differs in both its appearance and properties from the matters taken into the stomach, from which it


was prepared, and consequently cannot be the effect of mere solution, by which operation matter may be divided, but I presume can never be changed in its component parts. Again, chemistry has not yet taught us that any difference exists between the chyle of carnivorous and that of herbivorous animals, and from the most direct experiment we learn, that two animals of the same species being fed, the one on the matter of a muscular fibre, and the other on farinaceous matter, both afforded chyle in no respect different from each other.* Whereas, could chyle be produced by mere solution, it should surely differ in its properties, in proportion to the variety of matter from which it is prepared.

From these facts it appears, that the aid of some other operation is requisite to explain the formation of chyle. Perhaps several may be found necessary. In addition to solution, I do believe, that fermentation has likewise a very considerable effect. By it we know, bodies not only become decomposed and reduced to their elementary principles; but, by a recombination of those principles another substance is formed, differing materially from that from which it was obtained. It is by a similar decomposition and recombination of the elementary principles of food, together with its subsequent mixture with certain fluids of the primæ viæ, that I believe it becomes so far animalized and changed in its properties

* Fordyce on Food, p. 143.

perties as to form chyle. That by the combination of bodies another is formed, differing in its properties from either of those of which it is composed, is too true to be denied. Thus, if we combine with a proper proportion of hydrogen and carbon, a certain proportion of oxygen, we obtain sugar, a substance differing very essentially from either of the other three. The effect of solution not being that of a change of the component parts of food, it is clear that its only operation in digestion must be, that of expediting fermentation. This opinion, if we judge from analogy, I presume will be found correct. In similar circumstances out of the body, the more minutely any matter is divided which is capable of fermentation, the more speedily will that matter go through its several stages. To return to our simile of sugar. If we dissolve this substance in water, its particles may be so minutely divided as not to be perceptible in that fluid, yet by evaporation the same sugar may be obtained, not at all changed in its properties. But if we suffer sugar to ferment, the result will be very different. It will be resolved into its elementary principles, carbonic acid will be disengaged, and we will likewise obtain alcohol. Here then is a decomposition and recombination of its elementary principles. Sugar being composed of oxygen, hydrogen, and carbon, and these elements being separated by fermentation, are re-united to form these

two substances, to wit, the oxygene unites with a part of the carbone and is disengaged in the form of carbonic acid, while the remainder of the carbone is dissolved by the hydrogene and forms the alcohol. Thus we see the material difference in the effects of solution and fermentation.



SECTION IV.



WHY IT IS PRESUMABLE THAT FERMENTATION
TAKES PLACE IN DIGESTION.

WE have already considered digestion *a priori*; that is, we have investigated the causes which influence digestion. We have taken notice of the several properties of the matters which have the most material effect in this process. And we have likewise seen, that all the circumstances essential to fermentation, are possessed in an especial manner by food in the stomach. We have seen, for instance, that it is plentifully supplied with moisture, not only from our drinks, but even from our saliva and the fluids of the stomach itself; that it receives a sufficient quantity of air from our saliva, by which fluid it is enveloped and continually conveyed into the stomach; and lastly, that its situation is admirably adapted to be supplied with the necessary quantity

ity of heat; on all of which circumstances fermentation in a particular manner depends. We have also shewn, that saliva and the mucus of the primæ viæ, have a considerable tendency to promote this process.

Having proceeded thus far on our subject, it next becomes necessary, that we consider digestion *a posteriore*; or, in other words, that we attend to the effects produced by the digestion of food in the stomach. But in the first place, we shall say a few words on fermentation.

Fermentation is that great agent in nature, by which bodies are rendered totally different in their chemical properties, and which, from the variety of its products, has been long since divided into three several stages; to wit, the vinous, the acetous, and the putrefactive. From the first of these processes we obtain, *alcohol*; from the second, *vinegar*; and from the third, *ammoniac*; by which means we are able always to ascertain, the nature of whatever fermentation has taken place.

It will be recollected, that these several stages of fermentation are capable of taking place, entirely independent of each other. Whenever the saccharine principle of any matter predominates, the vinous fermentation will take place; when the mucilaginous is most abundant, it will undergo the acetous; and when a greater proportion of gluten is present, it will run immediately into the putrefactive

factive stage of fermentation. From this it appears, that on the several proportions of saccharine matter, of mucilage and of gluten, which any substance contains, depends the priority of the fermentation which will commence. Thus it is, that the vinous fermentation is capable of preceding the acetous, and *vice versa*. But they do not necessarily follow each other. Hence it is, that old and generous wines, in which the mucilaginous principle has been destroyed, are no longer capable of becoming acid, without the addition of a certain proportion of gummy matter.* Neither does milk afford a vinous spirit by its own spontaneous change, as in this case it loses its saccharine principle.

Different gases are disengaged, during the progress of these several stages of fermentation. The nature of these depend on the matter fermented. Thus, in the vinous and acetous stages of fermentation, carbonic acid is disengaged; while in the putrefactive, azote, carbonated hydrogen gas, sulphurated hydrogen gas, and phosphorated hydrogen gas, are all occasionally evolved.

We will now consider how far these gases, as well as the other products of fermentation, can be considered as being evolved in the *primæ viæ*.

In support of the first of these positions, we have that common fact, of our perceiving considerable eructations of air in affections of the stomach. Here

it

* Chaptal's Chem. Vol. III. page 262.

it may be said, that the disengagement of air is the consequence of disease. To this I answer, that the eructation I consider as such, but not the formation of air. In affections of the stomach from gout, the quantity of air disengaged is sometimes very great, yet we cannot suppose that it is formed by this affection. I regret that I cannot from my own experiments, say any thing relative to the nature of the air, which is evolved in the stomach during digestion. To place this matter, however, in as clear a light as I am able, I shall take the liberty of making a quotation from Plenck.* This gentleman informs us, that, “in a very healthy man, frozen to death by cold on a winter’s night, there was found a mixture of four kinds of air in the primæ viæ.

“Fixed air was found in the greatest quantity in the stomach, and but little in the small intestines.

“Vital air was contained chiefly in the stomach, and small intestines, and,

“Azote, and carbonated inflammable air, in the large intestines.”

To these I may add, that sulphurated hydrogen gas, and phosphorated hydrogen gas, have been proven to be disengaged *in crepitu*.

It appears from the works of Van Helmont, that he was the first person who suggested the idea of the presence of an acid in the stomach. His opinion

* Treat. on the human fluids, page 141.

nion has been ascribed to by Haller, who relates, that the acetous fermentation is very prevalent in the stomach; that an acid is spontaneously evolved before putrefaction, and even sometimes resists that process. He indeed mentions an acid being detected in the stomachs of ruminating and of omnivorous animals, in those of birds and even of carnivorous animals.*

Most persons have witnessed the presence of an acid in their stomachs. But here an objection arises. It has been said, that whenever an acid is present in the stomach it is morbid, and indicates the diseased state of that organ. When accompanied with eructations, I believe this to be the case; as the quantity then appears to be preternatural, and is attended with an inverted peristaltic motion of the stomach, which is certainly the effect of morbid action. But we must not infer from hence, that the presence of an acid in the stomach, is the effect of disease, as I hope to shew that it is detected in that organ, in the most sound and natural health.

This has been proven by Dr. Rush† many years ago. He has shewn us by several well directed experiments, that he always detected the presence of an acid in the contents of the stomach, when thrown up by an emetic, three hours after food was taken; but as it has been objected to his

H

experiments,

* *Elemen. Physiol.* Tom. VI. p. 140 and 141.

† *Inaug. Dissert. de Coctio, Cib. in Vent.*

experiments, that the acidity proceeded from the emetic tartar, which had been decomposed in the stomach, I shall endeavour to supersede this difficulty.

A gentleman in perfect health and capable of ruminating, dined on roasted beef, Irish potatoes and leavened bread. His drink was nothing but water. In four hours after, he brought up a portion of his dinner. It had an acid taste, and turned a blue vegetable substance of a red colour.

The same gentleman having dined on boiled mutton, cabbage and leavened bread, and having drunk water alone, as before, in four hours after, ruminated again. The portion of food brought up had an acid taste, and, as the last, turned a blue vegetable substance of a red colour.

These experiments were frequently repeated after having dined on different substances, and with uniform success. It was observed, that the acidity was not as perceptible to the taste in an hour or two after having dined, as it was in several hours after. These experiments may be readily repeated, as I do believe, that almost any person with a little trouble, may learn to ruminate.

Again, to determine whether an acid is evolved in the digestion of a cat, one was fed on boiled beef and Irish potatoes. In five hours after, it was strangled. Its abdomen was opened and its stomach taken out, having previously placed ligatures
on

on its two orifices. The food had become soft and pulpy, and there was very little of any kind of fluid in the stomach. Some of this pulpy mass being placed in a glass vessel and mixed with a little water, shewed evident marks of acidity. It very speedily restored the yellow colour of paper stained with rhubarb, after its having been turned brown by an alkali.

I am not ignorant of the assertion of Dr. For-
dyce, that in his experiments on dogs, cows, and
sheep, he could not find the least trace of acidity
in the duodenum;* neither could it have been ex-
pected otherwise, since, as has been already related,
the bile has the effect of neutralizing acids, and of
consequence as that fluid is almost always flowing
into the duodenum, the quantity of acid must be
unusually great to be detected in that intestine.

It is a prevalent opinion, that the acid which is
present in the stomach is the phosphoric, and con-
sequently that it is not the effect of fermentation.
But, as it is our duty not to admit either one
position or another, unless it is supported by facts
or experiments, I shall relate such of these as have
induced me to presume, that it is not the phospho-
ric acid which is usually found in the stomach.
And,

1st. The acid found in the stomach, does not
precipitate sugar of lead from its solution in water.

2d. " Being

* Vid. Treat. on Food. p. 150 and 151.

2d. "Being saturated with kali, that is, what
 "was formerly called fixed vegetable alkali, it
 "produces kali acetatum, formerly called regene-
 "rated tartar, or sal-diureticus."*

Desirous of knowing whether an acid is evolved
 in the digestion of animal substances, as well as
 in the former experiments, the gentleman, capable
 of ruminating, dined on roasted veal alone and
 drank water as usual. In four hours after, a por-
 tion of the contents of his stomach was brought up.
 It turned a blue vegetable substance of a red co-
 lour, and had an acid smell and taste. I confess my
 being at a loss in this case to determine, whether
 the acid was the effect of the digestion of the veal,
 or whether it proceeded from the remains of some
 former food. It was my intention to have ascer-
 tained this point, by repeating the experiment on
 the same person, after his having subsisted on ani-
 mal food for eight or ten days; but, as I have not
 now as many days previous to the delivering in
 of my piece, I shall be obliged of consequence to
 decline the idea. The following experiment how-
 ever, will at least adduce probability, in favour of
 the acid having been evolved by digestion.

Having placed two drachms of roasted veal in a
 glass vessel, and covered it completely with saliva, it
 spontaneously became acid, long previous to there
 being any signs of putrefaction taking place. This
 fact

* Fordyce on Food. p. 148.

fact is corroborated by the experiments of many authors of reputation. Haller takes notice of it.* Dr. Rush found that beef acquired an acid taste and smell, when exposed for two days to the heat of summer;† and Maclurg relates, that a mixture of mutton and water, passes through the acetous stage of fermentation before it putrefies.‡

All animal matters when mixed with fermentable vegetable substances, have a tendency to promote fermentation, as appears from a number of experiments instituted by Pringle, and who likewise adds, that “after such mixtures become sour they never return to a putrid state, but, on the contrary, grow more and more acid.”§ And yet, so far from our finding this ferment to be injurious to digestion, on the contrary, animal food seems to be the best adapted for the aliment of dyspeptic patients.

The publication of Dr. Wilson’s ingenious inaugural dissertation on digestion, has induced many persons to suppose, that every idea of fermentation taking place in digestion is unfounded. The Doctor submitted to distillation, the contents of his stomach, brought up by an emetic four hours after food was taken, with a view of ascertaining, whether he could detect the presence of spirit of wine. This he

* Vid. Element. Phisiol. Tom. VI. p. 316. † Vid. Inaug. Dif. p. 21.

‡ Treat. on the bile. p. 75.

§ Diseases of the Army, Appendix, paper 4, Experiment 28.

he could not, and from hence concludes, that fermentation does not take place in digestion. It will be seen that I have repeated the Doctor's experiment, and I am happy to add, with the same success; but I shall not draw the same conclusions.

Being in perfect health, I dined on corned beef, potatoes and leavened bread. My drink was water alone. In four hours after, I took twenty grains of ipecacuanha and brought up the contents of my stomach. It shewed evident marks of acidity. On submitting it to distillation, a transparent and limpid fluid came over into the receiver, which had a taste somewhat sweet, and an agreeable flavour; it had not the least appearance of spirit of wine, neither could I detect in it the presence of an acid. On examining however the *residuum*, which had not been evaporated to dryness, I was not a little surprised to find it still acid.

A dog was kept fasting for twenty-four hours. He was then fed for two days successively on animal food. Four hours and a half after taking his last meal, he was killed. The food in his stomach shewed evident marks of acidity. On submitting it to distillation, I obtained, as in the last experiment, a transparent and limpid fluid, somewhat sweet, with an agreeable flavour and in no respect different from that which I had obtained from the contents of my own stomach.

From the above experiments we learn, first, that an acid was produced in digestion as in the former experiments,

experiments, and that it was even found in the residuum after distillation; and secondly, that spirit of wine could not be obtained from the food by distillation; but by no means that fermentation did not take place. I do not suppose that the vinous fermentation in a healthy stomach, is ever so complete that spirit of wine is formed, this would be equally as incorrect as to say, that putrefaction takes place in that organ. Neither can we suppose, that spirit of wine can ever be obtained from any matter, in which the acetous stage of fermentation has been completed and is then present.

We have already seen that the acetous fermentation is capable of preceding the vinous; it is probable this may be the case in digestion; nor would it be by any means singular. In the preparation of *koumiss* from the milk of mares, a drink much in use among the Tartars, the acetous fermentation always precedes the vinous. But admitting that the vinous fermentation does have the priority, every circumstance, to which our food is exposed in the stomach, must tend to hurry it on so speedily to the acetous stage, as to prevent the former from being at all perceptible. The heat to which it is exposed in the stomach, is greater than that which is requisite for the vinous fermentation; and this excess of heat favours the acetous stage.

When speaking of the acid formed during digestion in the stomach, the word evolve has been used by

by many persons; whenever I have followed them in this respect, it will be recollected that I have always meant, that it was evolved by fermentation. I do not suppose, that the acid in the stomach is simply disengaged from our food, as it is from the common *sumach* when mixed with water, or any other substance in which it is very abundant. The quantity of acid contained in the food, on which we have experimented, is not so great that this could have been the case; neither can it be obtained from it out of the body by any other means, than by the assistance of fermentation.

It now remains with the reader to determine, whether or not my position is correct, that this acid is the effect of fermentation.

Fermentation out of the body, differs from that which takes place in a living stomach, in this, that the former is spontaneous; whereas, the latter is induced by all the numerous circumstances in the stomach, which tend to promote that process, and of consequence must be more speedy,

Does not the fact of digestion being more speedy while we are at rest, than during exercise, favour the theory of fermentation?

How shall we account for that warmth about the region of the stomach, so perceptible in some persons for several hours after death, unless we admit of fermentation?

Whether we shall ever be so successful as to imitate nature in digestion, as in many of her other operations,

operations, I am unable to decide; but, I flatter myself, time, that correct discerner of truth and error, will direct the attention of some more fortunate experimenter to this subject, and dissipate every doubt which may still involve it.

Having thus delivered my observations and experiments on digestion, in as concise a manner as my time would admit of, I shall now close this essay; but to do this without an acknowledgment to the several Medical Professors of this University, would be a breach of that duty which my feelings claim.

To you, gentlemen, at least this small tribute of my esteem is due. Permit me then to present you with the sincere acknowledgments of a pupil, for the many opportunities of improvement which your lectures have afforded him. With the assurance of my wishes, that you may continue to enjoy all the pleasure of success in the science of medicine, I now bid you, adieu.

F I N I S.

Med. Hist.

WZ

270

G566a

1800

41

